

An Information-Spectrum Approach to Classical and Quantum Hypothesis Testing for Simple Hypotheses

Hiroshi Nagaoka Masahito Hayashi

Abstract

The information-spectrum analysis made by Han for classical hypothesis testing for simple hypotheses is extended to a unifying framework including both classical and quantum hypothesis testing. The results are also applied to fixed-length source coding when loosening the normalizing condition for probability distributions and for quantum states. We establish general formulas for several quantities relating to the asymptotic optimality of tests/codes in terms of classical and quantum information spectra.

Index Terms

Information spectrum, Quantum hypothesis testing, Classical hypothesis testing, Fixed-length source coding, Optimal exponent

I. INTRODUCTION

One of the principal aims of information theory is to establish a link between two different kinds of quantities. One is an operational quantity which is defined as the optimal or limiting value of a concrete parameter such as code length, compression rate, transmission rate, convergence rate of error probabilities, etc. The other is an information quantity such as the entropy, divergence, mutual information, etc. Note that the latter, in its definition, is more abstract than the former, and the meaning of the latter is usually clarified by linking it to the former. In the so-called information spectrum method which first appeared in a series of joint papers of Han and Verdú (e.g., [1], [2]), the process of establishing such a link is intentionally divided into two parts by introducing a third kind of quantity — *information spectrum*, putting it between an operational quantity and an abstract information quantity. This setting allows us to pursue many problems of information theory in their most general forms; see [3] for the whole perspective of the method.

For instance, let us consider two sequences of random variables $\mathbf{X} = \{X^n\}_{n=1}^\infty$ and $\mathbf{Y} = \{Y^n\}_{n=1}^\infty$, where X^n and Y^n for each n are supposed to take values in a common discrete (finite or countable)¹ set $\mathcal{X}^n$ subject to probability distributions (mass functions) P_{X^n} and P_{Y^n} respectively. Note that $\mathcal{X}^n$ does not need to be the product set $\mathcal{X} \times \cdots \times \mathcal{X}$ of an $\mathcal{X}$, although the notation suggests that the product set is a representative example of $\mathcal{X}^n$. Han [3], [4] studied the hypothesis testing problem for the simple hypotheses consisting of the general processes $\mathbf{X}$ and $\mathbf{Y}$ by means of the information spectrum, which is the asymptotic behavior of the random variable $\frac{1}{n} \log \frac{P_{X^n}(X^n)}{P_{Y^n}(X^n)}$ (or $\frac{1}{n} \log \frac{P_{X^n}(Y^n)}{P_{Y^n}(Y^n)}$) in this case. He succeeded in representing several asymptotic characteristics of hypothesis testing in terms of the information spectrum with no or very few assumptions on the processes. The term ‘spectrum’ is intended to mean that the scope of the theory covers the general case when the probability distribution of $\frac{1}{n} \log \frac{P_{X^n}(X^n)}{P_{Y^n}(X^n)}$ does not necessarily get concentrated at a point, but may spread out, as $n \rightarrow \infty$.

The purpose of the present paper is to extend, complement and refine Han’s analysis of hypothesis testing from several viewpoints. The biggest motivation comes from the question of how to extend the analysis to quantum hypothesis testing. Following the above setting, we are naturally led to consider the problem of hypothesis testing for the simple hypotheses consisting of two sequences of quantum states $\vec{\rho} = \{\rho_n\}_{n=1}^\infty$ and $\vec{\sigma} = \{\sigma_n\}_{n=1}^\infty$, where ρ_n and σ_n are density operators on a common Hilbert space $\mathcal{H}_n$ for each n . However, it is by no means obvious whether a similar analysis to that of Han is applicable to the quantum setting. We show in this paper that it is actually possible to extend Han’s results by appropriately choosing a quantum analogue of the information spectrum so that both the classical and quantum cases are treated in a unifying framework. Although this does not mean that application to a special class of quantum processes such as i.i.d. (independent and identically distributed) ones immediately yields significant results, it seems to suggest a new approach to studying the quantum asymptotics and to elucidating a general principle underlying classical/quantum information theory.

Hiroshi Nagaoka is with Graduate School of Information Systems, The University of Electro-Communications, 1-5-1, Chofugaoka, Chofu-shi, Tokyo, 182-8585, Japan (e-mail: nagaoka@is.uec.ac.jp).

Masahito Hayashi was with the Laboratory for Mathematical Neuroscience, Brain Science Institute, RIKEN, 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan (e-mail: masahito@brain.riken.go.jp). He is now with ERATO-SORST Quantum Computation and Information Project, Japan Science and Technology Agency (JST), 201 Daini Hongo White Bldg, 5-28-3, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan (e-mail: masahito@qci.jst.go.jp), and Superrobust Computation Project, Information Science and Technology Strategic Core (21st Century COE by MEXT), Graduate School of Information Science and Technology, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.

¹ In this paper we only treat the discrete case to simplify the description when considering the classical hypothesis testing, although it is straightforward as pointed out in [3], [4] to extend the argument to the general case where $\{\mathcal{X}^n\}$ are arbitrary measurable spaces.

It should be noted that, even though the statements of our theorems are almost parallel to those for the classical setting, some of the proofs are essentially different from the original proofs of Han. The technique of information-spectrum slicing, which was effectively used in [3], [4], [6] to prove several important theorems, consists of a procedure of partitioning a set and does not straightforwardly apply to the quantum setting. We are thus forced to look for another idea for proofs. Fortunately, we have successfully found a way which does not need information-spectrum slicing and is applicable to the quantum setting. Moreover, the new proofs are much simpler than the original ones even in the classical case. This simplification is a byproduct of our attempt to pursue quantum extensions.

This paper also contains results such as those of Theorems 4 and 7 which improve the corresponding original theorems when applied to the classical setting. In addition, from the beginning, we treat generalized hypothesis testing in the sense of Han [3], [4], namely that the alternative hypothesis P_{Y^n} can be any nonnegative measure. This enables us to unify hypothesis testing and fixed-length source coding in a natural way.

This paper aims at presenting a unifying framework to treat the classical and quantum generalized hypothesis testing problem in the most general and simplest manner. After presenting the notation in section II, the concept of information spectrum is introduced in section III for both classical and quantum cases. In sections IV, VI and VII, various types of asymptotic bounds on the hypothesis testing problems for classical and quantum general processes are studied, basically following the problem settings and the notation given in [3], [4]. In section V we make some observations on Stein's lemma for classical and quantum i.i.d. processes in the light of the results of section IV. Applications to the classical fixed-length source coding are presented in section VIII, and concluding remarks are given in section IX.

II. A UNIFYING DESCRIPTION OF CLASSICAL AND QUANTUM GENERALIZED HYPOTHESIS TESTING

In this section we present a common language to treat classical and quantum hypothesis testing and fixed-length source coding in a unifying manner. We begin by considering the classical case. Suppose that we are given a sequence of discrete sets $\mathcal{X} = \{\mathcal{X}^n\}_{n=1}^\infty$, a sequence of probability measures $\vec{\rho} = \{\rho_n\}_{n=1}^\infty$ and a sequence of nonnegative (not necessarily probability) measures $\vec{\sigma} = \{\sigma_n\}_{n=1}^\infty$, which are represented by mass functions $\rho_n : \mathcal{X}^n \rightarrow [0, 1]$ with $\sum_{x \in \mathcal{X}^n} \rho_n(x) = 1$ and $\sigma_n : \mathcal{X}^n \rightarrow [0, \infty)$. In the usual hypothesis testing problem, both ρ_n and σ_n are probability measures denoted as $\rho_n = P_{X^n}$ and $\sigma_n = P_{Y^n}$. On the other hand, σ_n should be taken to be the counting measure on $\mathcal{X}^n$ when considering the source coding problem (see [3], [4] and section VIII below). For a function (random variable) $A : \mathcal{X}^n \rightarrow \mathbb{R}$, we write

$$\rho_n[A] = \sum_{x \in \mathcal{X}^n} \rho_n(x) A(x) \quad \text{and} \quad \sigma_n[A] = \sum_{x \in \mathcal{X}^n} \sigma_n(x) A(x).$$

Let $\mathcal{T}_n$ be the set of $[0, 1]$ -valued functions defined on $\mathcal{X}^n$. When both ρ_n and σ_n are probability measures, we regard an element T_n of $\mathcal{T}_n$ as a randomized test for the simple hypotheses $\{\rho_n, \sigma_n\}$ by interpreting $T_n(x) (\in [0, 1])$ as the probability of accepting the hypothesis ρ_n when the data x is observed. In particular, a deterministic test is an element of $\mathcal{T}_n$ taking values in $\{0, 1\}$, which is the characteristic function of the acceptance region $\{x \in \mathcal{X}^n \mid T_n(x) = 1\}$ for the hypothesis ρ_n . Depending on whether the true distribution is ρ_n or σ_n , the probability of accepting the hypothesis ρ_n turns out to be $\rho_n[T_n]$ or $\sigma_n[T_n]$, and the error probabilities of the first and second kinds are represented as

$$\alpha_n[T_n] \stackrel{\text{def}}{=} 1 - \rho_n[T_n] \quad \text{and} \quad \beta_n[T_n] \stackrel{\text{def}}{=} \sigma_n[T_n]. \quad (1)$$

In the general situation where σ_n is an arbitrary nonnegative measure, we still call elements of $\mathcal{T}_n$ tests and use the same notation as in (1). Letting I and 0 denote the constant functions on $\mathcal{X}^n$ such that $I(x) = 1$ and $0(x) = 0$ for all $x \in \mathcal{X}^n$, a test T_n is characterized as a function such that $0 \leq T_n \leq I$, where, and in the sequel, we write $A \leq B$ for functions A and B when $A(x) \leq B(x)$ for all x .

Let us turn to the quantum case. Suppose that a sequence of Hilbert spaces $\vec{\mathcal{H}} = \{\mathcal{H}_n\}_{n=1}^\infty$ is given. Let $\vec{\rho} = \{\rho_n\}_{n=1}^\infty$ be a sequence of density operators (i.e., nonnegative self-adjoint operators with trace one) on $\{\mathcal{H}_n\}$, and $\vec{\sigma} = \{\sigma_n\}_{n=1}^\infty$ be a sequence of bounded nonnegative self-adjoint (but not necessarily density or trace-class) operators on $\{\mathcal{H}_n\}$. For a bounded self-adjoint operator A on $\mathcal{H}_n$ we write

$$\rho_n[A] = \text{Tr}(\rho_n A) \quad \text{and} \quad \sigma_n[A] = \text{Tr}(\sigma_n A).$$

Let $\mathcal{T}_n$ be the set of self-adjoint operators T_n satisfying $0 \leq T_n \leq I$; i.e., both T_n and $I - T_n$ are nonnegative with I denoting the identity operator on $\mathcal{H}_n$. An element T_n of $\mathcal{T}_n$ can be considered to represent a $\{0, 1\}$ -valued measurement on $\mathcal{H}_n$ by identifying it with the POVM (positive operator-valued measure) $\{T_n(0), T_n(1)\} = \{T_n, I - T_n\}$, and is called a test for the hypotheses $\{\rho_n, \sigma_n\}$ with the interpretation that the measurement result 0 means the acceptance of the hypothesis ρ_n .

We define $\alpha_n[T_n]$ and $\beta_n[T_n]$ by the same equations as (1), which turn out to be the error probabilities of the first and second kinds when both ρ_n and σ_n are density operators.

We have thus reached a common setting to treat generalized hypothesis testing of classical and quantum systems for which sequences

$$\vec{\rho} = \{\rho_n\}_{n=1}^\infty, \quad \vec{\sigma} = \{\sigma_n\}_{n=1}^\infty, \quad \text{and} \quad \vec{\mathcal{T}} = \{\mathcal{T}_n\}_{n=1}^\infty$$

are given. Note that

$$0 = \rho_n[0] \leq \rho_n[T_n] \leq \rho_n[T'_n] \leq \rho_n[I] = 1 \quad (2)$$

and

$$0 = \sigma_n[0] \leq \sigma_n[T_n] \leq \sigma_n[T'_n] \leq \sigma_n[I] \leq \infty \quad (3)$$

always hold for any tests $T_n, T'_n \in \mathcal{T}_n$ such that $T_n \leq T'_n$. We shall work with this setting throughout this paper.

Remark 1: Readers who are familiar with the language of operator algebras may immediately extend the setting to a more general one in which we are given a sequence of a certain kind of $*$ -algebras $\vec{\mathcal{A}} = \{\mathcal{A}_n\}_{n=1}^\infty$ containing the identity elements I , a sequence of states $\vec{\rho} = \{\rho_n\}_{n=1}^\infty$ (linear functionals mapping nonnegative elements to nonnegative numbers and the identity elements to 1) and a sequence of positive weights $\vec{\sigma} = \{\sigma_n\}_{n=1}^\infty$ (linear functionals mapping nonnegative elements to nonnegative numbers or ∞), with defining $\mathcal{T}_n = \{T_n \in \mathcal{A}_n \mid 0 \leq T_n = T_n^* \leq I\}$. The classical case and the quantum case treated above correspond to $\mathcal{A}_n = L^\infty(\mathcal{X}^n)$ (the set of complex-valued bounded functions on $\mathcal{X}^n$) and $\mathcal{A}_n = \mathcal{B}(\mathcal{H}_n)$ (the set of bounded operators on $\mathcal{H}_n$) respectively.

The following notation is introduced in order to represent several variations of error exponents in a unifying manner. Given a sequence of tests $\vec{T} = \{T_n\}_{n=1}^\infty$ such that $T_n \in \mathcal{T}_n$ ($\forall n$), let

$$\begin{aligned} \eta_n[T_n] &\stackrel{\text{def}}{=} -\frac{1}{n} \log \alpha_n[T_n] = -\frac{1}{n} \log(1 - \rho[T_n]), \\ \zeta_n[T_n] &\stackrel{\text{def}}{=} -\frac{1}{n} \log \beta_n[T_n] = -\frac{1}{n} \log \sigma[T_n], \end{aligned}$$

and

$$\begin{aligned} \underline{\alpha}[\vec{T}] &\stackrel{\text{def}}{=} \liminf_{n \rightarrow \infty} \alpha_n[T_n], & \overline{\alpha}[\vec{T}] &\stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} \alpha_n[T_n], \\ \underline{\beta}[\vec{T}] &\stackrel{\text{def}}{=} \liminf_{n \rightarrow \infty} \beta_n[T_n], & \overline{\beta}[\vec{T}] &\stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} \beta_n[T_n], \\ \underline{\eta}[\vec{T}] &\stackrel{\text{def}}{=} \liminf_{n \rightarrow \infty} \eta_n[T_n], & \overline{\eta}[\vec{T}] &\stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} \eta_n[T_n], \\ \underline{\zeta}[\vec{T}] &\stackrel{\text{def}}{=} \liminf_{n \rightarrow \infty} \zeta_n[T_n], & \overline{\zeta}[\vec{T}] &\stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} \zeta_n[T_n]. \end{aligned}$$

When $\vec{T} = \{T_n\}$ is replaced with its *complement* $\vec{T}^c = \{T_n^c \stackrel{\text{def}}{=} I - T_n\}$, we add the superscript c to these symbols as $\alpha_n^c[T_n] = \alpha_n[T_n^c]$, $\eta_n^c[T_n] = \eta_n[T_n^c]$, $\underline{\zeta}^c[\vec{T}] = \underline{\zeta}[\vec{T}^c]$, etc.

III. INFORMATION SPECTRUM AND LIKELIHOOD TESTS

As mentioned in the introduction, the information spectrum for classical hypothesis testing is the asymptotic behavior of the random variable

$$Z_n \stackrel{\text{def}}{=} \frac{1}{n} \log \frac{\rho_n(X^n)}{\sigma_n(X^n)},$$

where X^n is supposed to be subject to the probability distribution ρ_n . Han [3], [4] called Z_n the divergence-density rate and derived several formulas for representing the asymptotic characteristics of the classical hypothesis testing problem in terms of the information spectrum. Now we are led to the following question; what is the quantum analogue of the information spectrum? At a first glance, it may seem to be natural to consider the quantum observable represented by the self-adjoint operator $\frac{1}{n}(\log \rho_n - \log \sigma_n)$ and its probability distribution under the quantum state ρ_n . Unfortunately, this line is not directly linked to the hypothesis testing problem. We give up seeking the quantum analogue of Z_n , but instead seek that of a likelihood test $S_n(a) : \mathcal{X}^n \rightarrow [0, 1]$ obeying

$$S_n(a)(x) = \begin{cases} 1 & \text{if } \frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} > a \\ 0 & \text{if } \frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} < a \end{cases} \quad (4)$$

where a is an arbitrary real number. Note that there is an ambiguity in this definition of $S_n(a)$ when some x satisfies $\frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} = a$, including two special cases where $S_n(a)$ are the deterministic tests with the acceptance regions

$$\begin{aligned} &\left\{ x \in \mathcal{X}^n \mid \frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} > a \right\} \\ &\text{and } \left\{ x \in \mathcal{X}^n \mid \frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} \geq a \right\}. \end{aligned}$$

In general $S_n(a)$ may be randomized with an arbitrary probability when the obtained data x satisfies $\frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} = a$. Denoting the characteristic functions of the sets $\{x \mid A(x) > c\}$ and $\{x \mid A(x) \geq c\}$ by $\{A > c\}$ and $\{A \geq c\}$ respectively, equation (4) is rewritten as

$$\left\{ \frac{1}{n} \log \frac{\rho_n}{\sigma_n} > a \right\} \leq S_n(a) \leq \left\{ \frac{1}{n} \log \frac{\rho_n}{\sigma_n} \geq a \right\}, \quad (5)$$

or equivalently as

$$\{\rho_n - e^{na}\sigma_n > 0\} \leq S_n(a) \leq \{\rho_n - e^{na}\sigma_n \geq 0\}. \quad (6)$$

The family of tests $\{S_n(a)\}_{a \in \mathbb{R}}$ characterizes the information spectrum by $\text{Prob}\{Z_n \geq a\} = \rho_n[S_n(a)]$.

In order to introduce the quantum analogue of $S_n(a)$, we need some preliminaries. For a self-adjoint operator A on a Hilbert space with the spectral decomposition² $A = \sum_i \lambda_i E_i$, where $\{\lambda_i\}$ are the eigenvalues and $\{E_i\}$ are the orthogonal projections onto the corresponding eigenspaces, we define

$$\{A \geq 0\} \stackrel{\text{def}}{=} \sum_{i: \lambda_i \geq 0} E_i \quad \text{and} \quad \{A > 0\} \stackrel{\text{def}}{=} \sum_{i: \lambda_i > 0} E_i.$$

These are the orthogonal projections onto the direct sum of eigenspaces corresponding to nonnegative and positive eigenvalues, respectively. The projections $\{A \leq 0\}$ and $\{A < 0\}$, or more generally $\{A \geq B\} = \{A - B \geq 0\}$, $\{A < B\} = \{A - B < 0\}$, etc., are defined similarly. We have

$$\text{Tr}(A\{A > 0\}) \geq 0, \quad (7)$$

and for any test T on $\mathcal{H}$

$$\text{Tr}(A\{A > 0\}) \geq \text{Tr}(AT). \quad (8)$$

The first inequality is obvious, while the second follows from $0 \leq T \leq I$ as

$$\begin{aligned} \text{Tr}(AT) &= \text{Tr}(A\{A > 0\}T) + \text{Tr}(A\{A \leq 0\}T) \\ &\leq \text{Tr}(A\{A > 0\}T) \\ &\leq \text{Tr}(A\{A > 0\}). \end{aligned}$$

Note that $\{A > 0\}$ in (7) and (8) can be replaced with $\{A \geq 0\}$ or, more generally, with any self-adjoint operator S satisfying $\{A > 0\} \leq S \leq \{A \geq 0\}$.

Now, in the quantum setting where a sequence of density operators $\vec{\rho} = \{\rho_n\}$ and that of bounded nonnegative self-adjoint operators $\vec{\sigma} = \{\sigma_n\}$ are given, let $S_n(a)$ be a self-adjoint operator satisfying the same equation as (6). Since $S_n(a)$ satisfies $0 \leq S_n(a) \leq I$, it is a test in our sense. Indeed, it is the quantum analogue of the likelihood test introduced by Holevo [7] and Helstrom [8] when σ_n is a density operator. Note that (6) is not equivalent to (5) in the quantum case unless ρ_n and σ_n commute. As in the classical case, there is an ambiguity in the definition of $S_n(a)$, including two special cases $S_n(a) = \{\rho_n - e^{na}\sigma_n \geq 0\}$ and $S_n(a) = \{\rho_n - e^{na}\sigma_n > 0\}$. Some quantities defined in the sequel may depend on a choice of $S_n(a)$ within (6), but this will not cause any essential difference in the theorems represented in terms of these quantities. We sometimes write $\{\rho_n - e^{na}\sigma_n \geq 0\}$ to mean $S_n(a)$, suggesting this ambiguity.

From (7) and (8) we have

$$(\rho_n - e^{na}\sigma_n)[S_n(a)] \geq 0, \quad (9)$$

and for any test T_n

$$(\rho_n - e^{na}\sigma_n)[S_n(a)] \geq (\rho_n - e^{na}\sigma_n)[T_n]. \quad (10)$$

In addition, letting $S_n^c(a) \stackrel{\text{def}}{=} I - S_n(a) = \{\rho_n - e^{na}\sigma_n \leq 0\}$ we have

$$(\rho_n - e^{na}\sigma_n)[S_n^c(a)] \leq 0, \quad (11)$$

$$(\rho_n - e^{na}\sigma_n)[S_n^c(a)] \leq (\rho_n - e^{na}\sigma_n)[T_n^c]. \quad (12)$$

These are rewritten as

$$\alpha_n(a) + e^{na}\beta_n(a) \leq 1, \quad (13)$$

$$\alpha_n(a) + e^{na}\beta_n(a) \leq \alpha_n[T_n] + e^{na}\beta_n[T_n], \quad (14)$$

and

$$\alpha_n(a) - e^{na}\beta_n^c(a) \leq 0, \quad (15)$$

$$\alpha_n(a) - e^{na}\beta_n^c(a) \leq \alpha_n[T_n] - e^{na}\beta_n^c[T_n], \quad (16)$$

²We assume here that A has discrete eigenvalues since it suffices for our main concern and simplifies the description, although the assumption is not essential.

where

$$\begin{aligned}\alpha_n(a) &\stackrel{\text{def}}{=} \alpha_n[S_n(a)] = \rho_n[\{\rho_n - e^{na}\sigma_n \leq 0\}], \\ \beta_n(a) &\stackrel{\text{def}}{=} \beta_n[S_n(a)] = \sigma_n[\{\rho_n - e^{na}\sigma_n \geq 0\}], \\ \beta_n^c(a) &\stackrel{\text{def}}{=} \beta_n[S_n^c(a)] = \sigma_n[\{\rho_n - e^{na}\sigma_n \leq 0\}].\end{aligned}$$

Needless to say, these properties also hold in the classical case. In particular, the inequality (14) in the classical case is the Neyman-Pearson lemma, whose quantum extension was given in [7], [8]. All the results in the later sections, including the classical ones obtained by Han, are derived only from the inequalities (13) through (16). This fact may be one of the most important findings of the present paper.

Let us see that $\alpha_n(a)$ ($\beta_n(a)$, resp.) is monotonically nondecreasing (nonincreasing, resp.) as a function of a ; i.e., if $a < b$ then

$$\alpha_n(a) \leq \alpha_n(b) \quad \text{and} \quad \beta_n(a) \geq \beta_n(b). \quad (17)$$

In the classical case, this is obvious because $\{x \mid \frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} > a\} \supset \{x \mid \frac{1}{n} \log \frac{\rho_n(x)}{\sigma_n(x)} \geq b\}$ if $a < b$. In order to show the monotonicity in the quantum case, we invoke (14) to yield

$$\begin{aligned}\alpha_n(a) + e^{na}\beta_n(a) &\leq \alpha_n(b) + e^{na}\beta_n(b), \\ \alpha_n(b) + e^{nb}\beta_n(b) &\leq \alpha_n(a) + e^{nb}\beta_n(a).\end{aligned}$$

These are rewritten as

$$e^{na}\{\beta_n(a) - \beta_n(b)\} \leq \alpha_n(b) - \alpha_n(a) \leq e^{nb}\{\beta_n(a) - \beta_n(b)\}, \quad (18)$$

which leads to (17). This monotonicity will be used implicitly throughout the later arguments.

Let

$$\begin{aligned}\eta_n(a) &\stackrel{\text{def}}{=} \eta_n[S_n(a)] = -\frac{1}{n} \log \alpha_n(a) \\ &= -\frac{1}{n} \log \rho_n[\{\rho_n - e^{na}\sigma_n \leq 0\}], \\ \zeta_n(a) &\stackrel{\text{def}}{=} \zeta_n[S_n(a)] = -\frac{1}{n} \log \beta_n(a) \\ &= -\frac{1}{n} \log \sigma_n[\{\rho_n - e^{na}\sigma_n \geq 0\}], \\ \zeta_n^c(a) &\stackrel{\text{def}}{=} \zeta_n^c[S_n(a)] = -\frac{1}{n} \log \beta_n^c(a) \\ &= -\frac{1}{n} \log \sigma_n[\{\rho_n - e^{na}\sigma_n \leq 0\}],\end{aligned}$$

and

$$\begin{aligned}\underline{\alpha}(a) &\stackrel{\text{def}}{=} \underline{\alpha}[\vec{S}(a)] = \liminf_{n \rightarrow \infty} \alpha_n(a), \\ \overline{\alpha}(a) &\stackrel{\text{def}}{=} \overline{\alpha}[\vec{S}(a)] = \limsup_{n \rightarrow \infty} \alpha_n(a), \\ \underline{\eta}(a) &\stackrel{\text{def}}{=} \underline{\eta}[\vec{S}(a)] = \liminf_{n \rightarrow \infty} \eta_n(a), \\ \underline{\zeta}(a) &\stackrel{\text{def}}{=} \underline{\zeta}[\vec{S}(a)] = \liminf_{n \rightarrow \infty} \zeta_n(a), \\ \overline{\zeta}^c(a) &\stackrel{\text{def}}{=} \overline{\zeta}^c[\vec{S}(a)] = \limsup_{n \rightarrow \infty} \zeta_n^c(a), \quad \text{etc.,}\end{aligned}$$

where $\vec{S}(a)$ denotes the sequence $\{S_n(a)\}_{n=1}^\infty$. Note that $\eta_n(a)$, $\zeta_n^c(a)$, $\underline{\eta}(a)$ and $\overline{\zeta}^c(a)$ are monotonically nonincreasing, while $\zeta_n(a)$ and $\underline{\zeta}(a)$ are monotonically nondecreasing. In addition, since (13) yields $\beta_n(a) \leq e^{-na}$, we have

$$\underline{\zeta}(a) \geq a. \quad (19)$$

IV. ASYMPTOTICS OF STEIN'S TYPE AND SPECTRAL DIVERGENCE RATES

In this section we treat the following quantities:

$$\begin{aligned}
B(\varepsilon | \vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \sup_{\vec{T}} \{ \zeta[\vec{T}] | \overline{\alpha}[\vec{T}] \leq \varepsilon \} \\
&= \sup \{ R | \exists \vec{T} \in \vec{\mathcal{T}}, \overline{\alpha}[\vec{T}] \leq \varepsilon \text{ and } \zeta[\vec{T}] \geq R \}, \\
B^\dagger(\varepsilon | \vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \sup_{\vec{T}} \{ \zeta[\vec{T}] | \underline{\alpha}[\vec{T}] < \varepsilon \} \\
&= \inf \{ R | \forall \vec{T} \in \vec{\mathcal{T}}, \text{ if } \zeta[\vec{T}] \geq R \text{ then } \underline{\alpha}[\vec{T}] \geq \varepsilon \},
\end{aligned}$$

where ε is a constant lying in the interval $[0, 1]$, and in particular

$$\begin{aligned}
B(\vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} B(0 | \vec{\rho} \| \vec{\sigma}) \\
&= \sup_{\vec{T}} \{ \zeta[\vec{T}] | \lim_{n \rightarrow \infty} \alpha_n[T_n] = 0 \} \\
&= \sup \{ R | \exists \vec{T}, \lim_{n \rightarrow \infty} \alpha_n[T_n] = 0 \text{ and } \zeta[\vec{T}] \geq R \}, \\
B^\dagger(\vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} B^\dagger(1 | \vec{\rho} \| \vec{\sigma}) \\
&= \sup_{\vec{T}} \{ \zeta[\vec{T}] | \underline{\alpha}[\vec{T}] < 1 \} \\
&= \inf \{ R | \forall \vec{T}, \text{ if } \zeta[\vec{T}] \geq R \text{ then } \lim_{n \rightarrow \infty} \alpha_n[T_n] = 1 \}.
\end{aligned}$$

As will be seen in the next section, these quantities are the main concern of Stein's lemma in the classical i.i.d. case. Note that we formally have $B(1 | \vec{\rho} \| \vec{\sigma}) = \infty$ and $B^\dagger(0 | \vec{\rho} \| \vec{\sigma}) = -\infty$, although they are of no importance. Obviously, for any $0 \leq \varepsilon_1 < \varepsilon_2 \leq 1$

$$\begin{aligned}
B(\vec{\rho} \| \vec{\sigma}) &\leq B(\varepsilon_1 | \vec{\rho} \| \vec{\sigma}) \leq B(\varepsilon_2 | \vec{\rho} \| \vec{\sigma}), \\
B^\dagger(\varepsilon_1 | \vec{\rho} \| \vec{\sigma}) &\leq B^\dagger(\varepsilon_2 | \vec{\rho} \| \vec{\sigma}) \leq B^\dagger(\vec{\rho} \| \vec{\sigma}),
\end{aligned}$$

and

$$B(\vec{\rho} \| \vec{\sigma}) \leq B(\varepsilon_1 | \vec{\rho} \| \vec{\sigma}) \leq B^\dagger(\varepsilon_2 | \vec{\rho} \| \vec{\sigma}) \leq B^\dagger(\vec{\rho} \| \vec{\sigma}).$$

In addition, $B(\varepsilon | \vec{\rho} \| \vec{\sigma})$ is right continuous for any $0 \leq \varepsilon < 1$ in the sense that

$$B(\varepsilon | \vec{\rho} \| \vec{\sigma}) = \max_{\vec{T}} \{ \zeta[\vec{T}] | \overline{\alpha}[\vec{T}] \leq \varepsilon \} = \inf_{\varepsilon' > \varepsilon} B(\varepsilon' | \vec{\rho} \| \vec{\sigma}). \quad (20)$$

To show this, let $\{\delta_k\}_{k=1}^\infty$ be an arbitrary sequence of positive numbers satisfying $\lim_{k \rightarrow \infty} \delta_k = 0$. Then for each k there exist a test $\vec{T}^{(k)} = \{T_n^{(k)}\}$ and a number n_k such that $\alpha_n[T_n^{(k)}] \leq \varepsilon + 2\delta_k$ and $\zeta_n[T_n^{(k)}] \geq B(\varepsilon + \delta_k | \vec{\rho} \| \vec{\sigma}) - \delta_k$ for all $n \geq n_k$. It is now easy to construct a test $\vec{T}$ such that $\overline{\alpha}[\vec{T}] \leq \varepsilon$ and $\zeta[\vec{T}] \geq \inf_{\varepsilon' > \varepsilon} B(\varepsilon' | \vec{\rho} \| \vec{\sigma})$, which proves (20). On the other hand, it is obvious that $B^\dagger(\varepsilon | \vec{\rho} \| \vec{\sigma})$ is left continuous for $0 < \varepsilon \leq 1$;

$$B^\dagger(\varepsilon | \vec{\rho} \| \vec{\sigma}) = \sup_{\varepsilon' < \varepsilon} B^\dagger(\varepsilon' | \vec{\rho} \| \vec{\sigma}). \quad (21)$$

Next, let

$$\begin{aligned}
\underline{D}(\varepsilon | \vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \sup \{ a | \overline{\alpha}(a) \leq \varepsilon \}, \\
\overline{D}(\varepsilon | \vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \sup \{ a | \underline{\alpha}(a) < \varepsilon \} = \inf \{ a | \underline{\alpha}(a) \geq \varepsilon \}
\end{aligned}$$

for $0 \leq \varepsilon \leq 1$, and

$$\begin{aligned}
\underline{D}(\vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \underline{D}(0 | \vec{\rho} \| \vec{\sigma}) = \sup \{ a | \lim_{n \rightarrow \infty} \alpha_n(a) = 0 \}, \\
\overline{D}(\vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \overline{D}(1 | \vec{\rho} \| \vec{\sigma}) = \inf \{ a | \lim_{n \rightarrow \infty} \alpha_n(a) = 1 \}.
\end{aligned}$$

It should be noted that in the classical case when $\rho_n = P_{X^n}$ and $\sigma_n = P_{Y^n}$ we have

$$\underline{D}(\vec{\rho} \| \vec{\sigma}) = \text{p-lim inf}_{n \rightarrow \infty} \frac{1}{n} \log \frac{P_{X^n}(X^n)}{P_{Y^n}(X^n)}$$

and

$$\overline{D}(\vec{\rho} \| \vec{\sigma}) = \text{p-lim sup}_{n \rightarrow \infty} \frac{1}{n} \log \frac{P_{X^n}(X^n)}{P_{Y^n}(X^n)},$$

where $p\text{-lim inf}$ and $p\text{-lim sup}$ are the liminf and limsup in probability:

$$\begin{aligned} p\text{-lim inf}_{n \rightarrow \infty} A_n &= \sup \{a \mid \lim_{n \rightarrow \infty} \text{Prob}\{A_n \geq a\} = 1\}, \\ p\text{-lim sup}_{n \rightarrow \infty} A_n &= \inf \{a \mid \lim_{n \rightarrow \infty} \text{Prob}\{A_n \leq a\} = 0\}. \end{aligned}$$

Actually, $\underline{D}(\mathbf{X} \parallel \mathbf{Y})$ and $\overline{D}(\mathbf{X} \parallel \mathbf{Y})$ were introduced in [3], [4] by these expressions and called the spectral sup- and inf-divergence rates between $\mathbf{X}$ and $\mathbf{Y}$.

It is clear that for any $0 \leq \varepsilon_1 < \varepsilon_2 \leq 1$

$$\begin{aligned} \underline{D}(\vec{\rho} \parallel \vec{\sigma}) &\leq \underline{D}(\varepsilon_1 \mid \vec{\rho} \parallel \vec{\sigma}) \leq \underline{D}(\varepsilon_2 \mid \vec{\rho} \parallel \vec{\sigma}), \\ \overline{D}(\varepsilon_1 \mid \vec{\rho} \parallel \vec{\sigma}) &\leq \overline{D}(\varepsilon_2 \mid \vec{\rho} \parallel \vec{\sigma}) \leq \overline{D}(\vec{\rho} \parallel \vec{\sigma}), \end{aligned}$$

and

$$\underline{D}(\vec{\rho} \parallel \vec{\sigma}) \leq \underline{D}(\varepsilon_1 \mid \vec{\rho} \parallel \vec{\sigma}) \leq \overline{D}(\varepsilon_2 \mid \vec{\rho} \parallel \vec{\sigma}) \leq \overline{D}(\vec{\rho} \parallel \vec{\sigma}).$$

In addition, when $\vec{\sigma} = \{\sigma_n\}$ consists of states (probability distributions in the classical case and density operators in the quantum case) as well as $\vec{\rho} = \{\rho_n\}$, it follows from (15) that

$$\alpha_n(a) \leq e^{na}(1 - \beta_n(a)) \leq e^{na}. \quad (22)$$

Hence we have $\lim_{n \rightarrow \infty} \alpha_n(a) = 0$ for any $a < 0$, which leads to

$$\underline{D}(\vec{\rho} \parallel \vec{\sigma}) \geq 0. \quad (23)$$

On the other hand, $\underline{D}(\vec{\rho} \parallel \vec{\sigma})$ and $\overline{D}(\vec{\rho} \parallel \vec{\sigma})$ may be negative when $\{\sigma_n\}$ are not states.

Theorem 1: For every $\varepsilon \in [0, 1]$

$$B(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) = \underline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}), \quad (24)$$

$$B^\dagger(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) = \overline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}). \quad (25)$$

In particular, we have

$$B(\vec{\rho} \parallel \vec{\sigma}) = \underline{D}(\vec{\rho} \parallel \vec{\sigma}), \quad (26)$$

$$B^\dagger(\vec{\rho} \parallel \vec{\sigma}) = \overline{D}(\vec{\rho} \parallel \vec{\sigma}). \quad (27)$$

Proof: Recalling that $\alpha_n(a) = \alpha_n[S_n(a)]$ and using equation (19), we have

$$\begin{aligned} B(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) &\geq \sup_a \{\zeta(a) \mid \overline{\alpha}(a) \leq \varepsilon\} \\ &\geq \sup \{a \mid \overline{\alpha}(a) \leq \varepsilon\} = \underline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}). \end{aligned}$$

To show the converse inequality, suppose that a test $\vec{T}$ and a real number a satisfy

$$\overline{\alpha}[\vec{T}] \leq \varepsilon < \overline{\alpha}(a).$$

Note that we can assume with no loss of generality the existence of such an a , or equivalently the finiteness of $\underline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma})$, since the inequality is trivial otherwise. Then there exists a positive δ for which $\alpha_n(a) - \alpha_n[T_n] \geq \delta$ holds for infinitely many n 's. Using equation (14) we have

$$\beta_n[T_n] \geq e^{-na} \{\alpha_n(a) - \alpha_n[T_n]\} + \beta_n(a) \geq e^{-na} \delta$$

for these n 's, which implies that $\zeta[\vec{T}] \leq a$. This proves

$$B(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) \leq \inf \{a \mid \overline{\alpha}(a) > \varepsilon\} = \underline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}).$$

Equation (24) has thus been verified. We can also prove equation (25) almost in the same way. ■

Remark 2: Equation (24) for the classical case was obtained by Han [3] as a slight modification of a result by Chen [9]. Equation (26) was also described in [3], giving credit to Verdú [10] for the original reference. As was mentioned in [3] and is now obvious from (26) and (27), the equality $\underline{D}(\vec{\rho} \parallel \vec{\sigma}) = \overline{D}(\vec{\rho} \parallel \vec{\sigma})$ is necessary and sufficient for the so-called strong converse property to hold in the sense that $\alpha_n[T_n]$ converges to 1 for any test $\vec{T}$ satisfying $\zeta[\vec{T}] > B(\vec{\rho} \parallel \vec{\sigma})$.

V. STEIN'S LEMMA IN THE CLASSICAL AND QUANTUM I.I.D. CASE

In the classical i.i.d. case when $\rho_n(x_1, \dots, x_n) = \rho(x_1) \cdots \rho(x_n)$ and $\sigma_n(x_1, \dots, x_n) = \sigma(x_1) \cdots \sigma(x_n)$, Stein's lemma (e.g., [11], [12], [13]) claims that

$$B(\varepsilon_1 | \vec{\rho} \| \vec{\sigma}) = B^\dagger(\varepsilon_2 | \vec{\rho} \| \vec{\sigma}) = D(\rho \| \sigma) \quad (28)$$

for $0 \leq \forall \varepsilon_1 < 1$, $0 < \forall \varepsilon_2 \leq 1$, where $D(\rho \| \sigma)$ is the Kullback-Leibler divergence: $D(\rho \| \sigma) = \sum_x \rho(x) \log \frac{\rho(x)}{\sigma(x)}$. A standard proof of the lemma uses a similar argument to the proof of Theorem 1 to reduce (28) to

$$\underline{D}(\vec{\rho} \| \vec{\sigma}) = \overline{D}(\vec{\rho} \| \vec{\sigma}) = D(\rho \| \sigma), \quad (29)$$

which is equivalent to

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\rho_n(X_1, \dots, X_n)}{\sigma_n(X_1, \dots, X_n)} = D(\rho \| \sigma) \quad \text{in probability,}$$

where $X_1, \dots, X_n$ are random variables obeying the probability distribution ρ_n . Now this is a direct consequence of the weak law of large numbers.

Let us turn to the quantum i.i.d. case when $\rho_n = \rho^{\otimes n}$ and $\sigma_n = \sigma^{\otimes n}$, where ρ and σ are density operators on a Hilbert space $\mathcal{H}$, and define the quantum relative entropy by $D(\rho \| \sigma) = \text{Tr} [\rho(\log \rho - \log \sigma)]$ (e.g., [14]). The achievability part

$$B(\varepsilon | \vec{\rho} \| \vec{\sigma}) \geq D(\rho \| \sigma) \quad \text{for } 0 < \forall \varepsilon < 1, \quad (30)$$

which is equivalent to $B(\vec{\rho} \| \vec{\sigma}) \geq D(\rho \| \sigma)$ by (20), was first proved by Hiai and Petz [15]. They showed the existence of a sequence of POVMs $\vec{M} = \{M^{(n)}\}$ on $\{\mathcal{H}^{\otimes n}\}$ satisfying

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{M^{(n)}}(\rho^{\otimes n} \| \sigma^{\otimes n}) \geq D(\rho \| \sigma), \quad (31)$$

where $D_{M^{(n)}}(\rho^{\otimes n} \| \sigma^{\otimes n})$ denotes the Kullback-Leibler divergence between the probability distributions $P_{\rho^{\otimes n}}^{M^{(n)}}(\cdot) = \text{Tr}(\rho^{\otimes n} M^{(n)}(\cdot))$ and $P_{\sigma^{\otimes n}}^{M^{(n)}}(\cdot) = \text{Tr}(\sigma^{\otimes n} M^{(n)}(\cdot))$. Since $nD(\rho \| \sigma) = D(\rho^{\otimes n} \| \sigma^{\otimes n}) \geq D_{M^{(n)}}(\rho^{\otimes n} \| \sigma^{\otimes n})$ follows from the monotonicity of relative entropy, this leads to

$$D(\rho \| \sigma) = \lim_{n \rightarrow \infty} \frac{1}{n} \sup_{M^{(n)}} D_{M^{(n)}}(\rho^{\otimes n} \| \sigma^{\otimes n}), \quad (32)$$

which is often referred to as the Hiai-Petz theorem. Now it is easy to see that combination of (31) and the direct part of the classical Stein's lemma leads to (30) as is shown in [15]. Hayashi [16] gave another construction of $\{M^{(n)}\}$ satisfying (31) based on a representation-theoretic consideration³. Inequality (30) can also be proved more directly, not by way of (31), in several different ways as shown in [17], [18] and Remark 20 of [19], the last of which also appears in Sec. 3.6 of [20]. Note that these proofs conversely yield the existence of $\vec{M} = \{M^{(n)}\}$ achieving (31) with the help of (35) below.

On the other hand, the converse part $B^\dagger(\vec{\rho} \| \vec{\sigma}) \leq D(\rho \| \sigma)$ was first shown in [21] by combining the quantum Neyman-Pearson lemma (14) with the inequality⁴

$$\text{Tr}(\rho^{\otimes n} \{\rho^{\otimes n} - e^{na} \sigma^{\otimes n} \geq 0\}) \leq e^{-n\{as - \psi(\theta)\}} \quad (33)$$

for $0 \leq \forall \theta \leq 1$, where $\psi(\theta) \stackrel{\text{def}}{=} \log \text{Tr}(\rho^{1+\theta} \sigma^{-\theta})$. A simpler proof was given in [22].

The quantum Stein's lemma has thus been established in the same form as (28). In the quantum case, (29) is not a ground of (28), as at present we do not have a quantum version of the law of large numbers which directly applies to (29) (even though only the inequality $\overline{D}(\vec{\rho} \| \vec{\sigma}) \leq D(\rho \| \sigma)$ follows immediately from (33)). Instead, (29) should be regarded as a consequence of (28). So we restate it as a theorem.

Theorem 2: For arbitrary density operators ρ and σ on a Hilbert space, we have $\underline{D}(\vec{\rho} \| \vec{\sigma}) = \overline{D}(\vec{\rho} \| \vec{\sigma}) = D(\rho \| \sigma)$ by letting $\vec{\rho} = \{\rho^{\otimes n}\}$ and $\vec{\sigma} = \{\sigma^{\otimes n}\}$; in other words,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \text{Tr}(\rho^{\otimes n} \{\rho^{\otimes n} - e^{na} \sigma^{\otimes n} \geq 0\}) \\ &= \begin{cases} 1 & \text{if } a < D(\rho \| \sigma), \\ 0 & \text{if } a > D(\rho \| \sigma). \end{cases} \end{aligned} \quad (34)$$

Example 1: Let us numerically illustrate this theorem for

$$\rho = \begin{bmatrix} 0.75 & 0.35 \\ 0.35 & 0.25 \end{bmatrix} \quad \text{and} \quad \sigma = \begin{bmatrix} 0.9 & 0 \\ 0 & 0.1 \end{bmatrix},$$

which are density operators (matrices) on $\mathcal{H} = \mathbb{C}^2$. The relative entropy in this case is $D(\rho \| \sigma) = 0.4013 \cdots$. The graph of the function $g_n(a) \stackrel{\text{def}}{=} \text{Tr}(\rho^{\otimes n} \{\rho^{\otimes n} - e^{na} \sigma^{\otimes n} > 0\})$ for $n = 5, 15$ and 50 is shown in Fig.1, where we can see that the

³More precisely, the papers [15] and [16] showed different theorems, both of which include (31) as a special case; see [16] for details.

⁴As a consequence of Eq. (2.63) in [20], the inequality of (33) turns out to be true for $\forall \theta \geq 0$.

slope of the graph around $a = D(\rho \parallel \sigma)$ gets steeper with increase of n , as equation (34) suggests. It is noted that drawing the graph requires computing the spectral decomposition of the $2^n \times 2^n$ matrix $\rho^{\otimes n} - e^{na} \sigma^{\otimes n}$ for each a , which is too large to apply a direct method when $n = 50$. We have applied the theory of irreducible decomposition of the algebra generated by $\{A^{\otimes n} \mid A \in \mathbb{C}^{2 \times 2}\}$ based on the observation made in [16], [17], which reduces the problem to finding the spectral decompositions of $\lceil (n+1)/2 \rceil$ matrices whose sizes are at most $(n+1) \times (n+1)$. Details of the algorithm will be reported elsewhere.




Fig. 1. The graph of $g_n(a)$

Let us make some observations on the quantum Neyman-Pearson test $S_n(a)$ in connection with (31). We begin by considering the general situation where $\vec{\rho} = \{\rho_n\}$ and $\vec{\sigma} = \{\sigma_n\}$ are arbitrarily given. For a sequence of tests $\vec{T} = \{T_n\}$, let $D_{T_n}(\rho_n \parallel \sigma_n)$ denote the Kullback-Leibler divergence of the resulting probability distributions $(\rho_n[T_n], 1 - \rho_n[T_n])$ and $(\sigma_n[T_n], 1 - \sigma_n[T_n])$. Then we have

$$\begin{aligned} D_{T_n}(\rho_n \parallel \sigma_n) &= -h(\rho_n[T_n]) - \rho_n[T_n] \log \sigma_n[T_n] \\ &\quad - (1 - \rho_n[T_n]) \log(1 - \sigma_n[T_n]) \\ &\geq -\log 2 - \rho_n[T_n] \log \sigma_n[T_n], \end{aligned}$$

where h is the binary entropy function; $h(t) \stackrel{\text{def}}{=} -t \log t - (1-t) \log(1-t)$. This proves that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{T_n}(\rho_n \parallel \sigma_n) \geq \underline{\zeta}[\vec{T}] \quad \text{if} \quad \lim_{n \rightarrow \infty} \alpha_n[T_n] = 0. \quad (35)$$

In particular, we have

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{S_n(a)}(\rho_n \parallel \sigma_n) \geq a \quad \text{if} \quad a < \underline{D}(\vec{\rho} \parallel \vec{\sigma}), \quad (36)$$

where we have used (19). Applying this to the i.i.d. case where $\rho_n = \rho^{\otimes n}$ and $\sigma_n = \sigma^{\otimes n}$, and recalling Theorem 2, we have

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{S_n(a)}(\rho^{\otimes n} \parallel \sigma^{\otimes n}) \geq a \quad \text{if} \quad a < D(\rho \parallel \sigma).$$

Therefore, if a number sequence $\{a_n\}$ is chosen so that a_n converges to $D(\rho \parallel \sigma)$ monotonically from below with a sufficient slow speed, then

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{S_n(a_n)}(\rho^{\otimes n} \parallel \sigma^{\otimes n}) \geq D(\rho \parallel \sigma), \quad (37)$$

which gives an example of (31).

Remark 3: From (36) and the monotonicity of quantum relative entropy, we obtain the general inequality

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D(\rho_n \parallel \sigma_n) \geq \underline{D}(\vec{\rho} \parallel \vec{\sigma}),$$

which has also appeared in [19].

Remark 4: Recently an extension of quantum Stein's lemma was reported under the name of "a quantum version of Sanov's theorem" [23]. Relating to this work, let us make some remarks on the relation between Stein's lemma and Sanov's theorem (see e.g. [13]). These theorems are similar in that both of them represent the convergence rates of some probabilities in terms of relative entropy. Moreover, they are closely related to each other in their logical derivations. Nevertheless, we should note that they have their respective roles in different contexts in general; Stein's lemma is about hypothesis testing and Sanov's theorem is a fundamental theorem in large deviation theory for empirical distribution. We also note that distinction of their roles is indispensable for precise understanding of both the significance of the Neyman-Pearson lemma and that of empirical distributions. Even though the result of [23] has a certain significance from a viewpoint of hypothesis testing, its formulation does not precisely correspond to that of Sanov's theorem in the classical case. Finding a meaningful and useful quantum extension of Sanov's theorem is still a challenging open problem. See also Remark 8 below.

VI. TRADEOFF BETWEEN THE EXPONENTS OF THE FIRST AND SECOND KIND ERROR PROBABILITIES

In order to properly evaluate the tradeoff between the error exponents of the first and second kinds for the classical hypothesis testing problem, Han [3], [4] introduced the following quantity:

$$\begin{aligned} B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) & \stackrel{\text{def}}{=} \sup \{ R \mid \exists \vec{T} \in \vec{\mathcal{T}}, \underline{\eta}[\vec{T}] \geq r \text{ and } \underline{\zeta}[\vec{T}] \geq R \} \\ & = \sup_{\vec{T}} \{ \underline{\zeta}[\vec{T}] \mid \underline{\eta}[\vec{T}] \geq r \}, \end{aligned} \quad (38)$$

where the subscript e is intended to mean that the quantity concerns exponents. Roughly speaking, the second kind error probability optimally tends to 0 with the rate $\beta_n[T_n] \approx e^{-nB_e(r \mid \vec{\rho} \parallel \vec{\sigma})}$ when the first error probability is required to tend to 0 with $\alpha_n[T_n] \approx e^{-nr}$ or faster. The same definition is applied to our setting including generalized and quantum hypothesis testing problem. We shall give some characterizations to this quantity in the sequel, extending the formula obtained by Han. The following lemma will play an essential role.

Lemma 1: For any real number a and any sequence of tests $\vec{T}$ we have

$$\underline{\zeta}(a) \geq \min \{ \underline{\zeta}[\vec{T}], a + \underline{\eta}[\vec{T}] \}, \quad \text{and} \quad (39)$$

$$\underline{\eta}(a) \geq \min \{ \underline{\eta}[\vec{T}], -a + \underline{\zeta}[\vec{T}] \}. \quad (40)$$

In particular, for any $a < b$

$$\underline{\zeta}(a) \geq a + \underline{\eta}(b) \quad \text{if} \quad \underline{\zeta}(a) < \underline{\zeta}(b), \quad \text{and} \quad (41)$$

$$\underline{\eta}(b) \geq -b + \underline{\zeta}(a) \quad \text{if} \quad \underline{\eta}(a) > \underline{\eta}(b). \quad (42)$$

Proof: We have

$$\begin{aligned} & \underline{\zeta}(a) \\ & = - \limsup_{n \rightarrow \infty} \frac{1}{n} \log \beta_n(a) \\ & \stackrel{(i)}{=} - \max \left\{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log \beta_n[T_n], \limsup_{n \rightarrow \infty} \frac{1}{n} \log (\beta_n(a) - \beta_n[T_n]) \right\} \\ & = \min \left\{ \underline{\zeta}[\vec{T}], \liminf_{n \rightarrow \infty} -\frac{1}{n} \log [\beta_n(a) - \beta_n[T_n]] \right\} \\ & \stackrel{(ii)}{\geq} \min \{ \underline{\zeta}[\vec{T}], a + \underline{\eta}[\vec{T}] \}, \end{aligned}$$

where the equality $\stackrel{(i)}{=}$ follows from the formula

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{n} \log (x_n + y_n) \\ & = \max \left\{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log x_n, \limsup_{n \rightarrow \infty} \frac{1}{n} \log y_n \right\} \end{aligned} \quad (43)$$

which is valid for any sequences of positive numbers $\{x_n\}, \{y_n\}$, and the inequality $\stackrel{(ii)}{\geq}$ follows from (14). The inequality (39) is thus proved. The proof of (40) is similar and omitted. $\blacksquare$

Theorem 3: For any $r \geq 0$,

$$\begin{aligned} B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) & = \sup_a \{ \underline{\zeta}(a) \mid \underline{\eta}(a) \geq r \} \\ & = \inf_a \{ a + \underline{\eta}(a) \mid \underline{\eta}(a) < r \} \\ & = \underline{\zeta}(a_0 - 0) = a_0 + \underline{\eta}(a_0 + 0), \end{aligned}$$

where

$$a_0(\in \mathbb{R} \cup \{-\infty, \infty\}) \\ \stackrel{\text{def}}{=} \sup \{a' \mid \exists a, \underline{\eta}(a) \geq r \text{ and } \underline{\zeta}(a') = \underline{\zeta}(a)\}.$$

Remark 5: Throughout the paper we use the notation

$$f(a+0) = \lim_{\varepsilon \downarrow 0} f(a+\varepsilon), \quad f(a-0) = \lim_{\varepsilon \downarrow 0} f(a-\varepsilon), \\ f(\infty) = \lim_{a \rightarrow \infty} f(a), \quad f(-\infty) = \lim_{a \rightarrow -\infty} f(a)$$

for a monotone function $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{-\infty, \infty\}$.

Remark 6: The formula given in the above theorem is valid even in the ‘singular’ case when the set $\{a \mid \underline{\eta}(a) \geq r\}$ is empty or the entire real line $\mathbb{R}$. When the set is empty, we have $B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) = -\infty$, although this does not occur in the case when σ_n are states (i.e., $\sigma_n[I] = 1$). When the set is $\mathbb{R}$, we have $B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) = \infty$.

Proof of Theorem 3: Since $\underline{\eta}(a) = \underline{\eta}[\vec{S}(a)]$ and $\underline{\zeta}(a) = \underline{\zeta}[\vec{S}(a)]$, it is immediate from the definition (38) of $B_e(r \mid \vec{\rho} \parallel \vec{\sigma})$ that

$$B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) \geq \sup_a \{ \underline{\zeta}(a) \mid \underline{\eta}(a) \geq r \}. \quad (44)$$

Also, we immediately have

$$\begin{aligned} & \sup_a \{ \underline{\zeta}(a) \mid \underline{\eta}(a) \geq r \} \\ &= \sup_{a'} \{ \underline{\zeta}(a') \mid \exists a, \underline{\eta}(a) \geq r \text{ and } \underline{\zeta}(a') = \underline{\zeta}(a) \} \\ &\geq \underline{\zeta}(a_0 - 0) \end{aligned} \quad (45)$$

and

$$a_0 + \underline{\eta}(a_0 + 0) \geq \inf_a \{ a + \underline{\eta}(a) \mid \underline{\eta}(a) < r \}. \quad (46)$$

Next, for any sequence of tests $\vec{T} = \{T_n\}$ satisfying $\underline{\eta}[\vec{T}] \geq r$ and for any number a satisfying $\underline{\eta}(a) < r$, it follows from (40) that $\underline{\zeta}[\vec{T}] \leq a + \underline{\eta}(a)$, which proves

$$B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) \leq \inf_a \{ a + \underline{\eta}(a) \mid \underline{\eta}(a) < r \}. \quad (47)$$

Finally, let us show that

$$\underline{\zeta}(a_0 - 0) \geq a_0 + \underline{\eta}(a_0 + 0), \quad (48)$$

which, combined with (44) through (47), completes the proof of the theorem. Assume first that the set $\{a \mid \underline{\eta}(a) \geq r\}$ is not empty nor $\mathbb{R}$. Then, letting b be an arbitrary number satisfying $\underline{\eta}(b) < r$, it follows from (42) that

$$\sup_a \{ \underline{\zeta}(a) \mid \underline{\eta}(a) \geq r \} \leq b + \underline{\eta}(b) < b + r < \infty.$$

Therefore, invoking that $\lim_{a \rightarrow \infty} \underline{\zeta}(a) = \infty$ follows from (19), we see that a_0 is not ∞ nor $-\infty$. Now, for an arbitrary $\varepsilon > 0$, we have $\underline{\zeta}(a_0 - \varepsilon) < \underline{\zeta}(a_0 + \varepsilon)$ by the definition of a_0 . Hence, from (41) of Lemma 1 we have

$$\underline{\zeta}(a_0 - \varepsilon) \geq a_0 - \varepsilon + \underline{\eta}(a_0 + \varepsilon),$$

which leads to (48). When $\{a \mid \underline{\eta}(a) \geq r\} = \emptyset$, on the other hand, we have $a_0 = -\infty$, and (48) is obvious since the right-hand side is $-\infty$. When $\{a \mid \underline{\eta}(a) \geq r\} = \mathbb{R}$, we have $a_0 = \infty$ and, again, (48) is obvious since the left-hand side is ∞ . ■

Remark 7: The formula $B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) = \inf_a \{ a + \underline{\eta}(a) \mid \underline{\eta}(a) < r \}$ for the classical hypothesis testing was derived by Han [3], [4], using a more complicated argument based on the technique of information-spectrum slicing. Note that this technique consists of a procedure of partitioning a set and does not straightforwardly apply to the quantum setting. Theorem 3 of [9] was also intended to give a general formula for the tradeoff between the error exponents of the first and second kinds, but its proof contains a gap, and the theorem does not apply to the general case where $\underline{\eta}(a)$ and $\underline{\zeta}(a)$ may be discontinuous functions; see Example 3.6 of [4].

Example 2: Let us consider the case when $\vec{\rho}$ and $\vec{\sigma}$ consist of pure quantum states of the form $\rho_n = |\psi_n\rangle\langle\psi_n|$ and $\sigma_n = |\varphi_n\rangle\langle\varphi_n|$, where ψ_n and φ_n are unit vectors in a Hilbert space $\mathcal{H}_n$ for every n , and let $\delta_n \stackrel{\text{def}}{=} \text{Tr}(\rho_n \sigma_n) = |\langle\psi_n|\varphi_n\rangle|^2$. In order to treat the hypothesis testing problem for this situation, we can assume with no loss of generality that

$$\psi_n = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \varphi_n = \begin{bmatrix} \sqrt{\delta_n} \\ \sqrt{1-\delta_n} \end{bmatrix},$$

for which we have the following spectral decomposition:

$$\rho_n - e^{na}\sigma_n = \lambda_1 E_1 + \lambda_2 E_2,$$

where

$$\lambda_1 = \frac{1 - e^{na}}{2} + r, \quad \lambda_2 = \frac{1 - e^{na}}{2} - r,$$

$$E_1 = \frac{1}{2} \begin{bmatrix} 1 + \frac{1 + e^{na} - 2e^{na}\delta_n}{2r} & -\frac{e^{na}\sqrt{\delta_n(1-\delta_n)}}{r} \\ -\frac{e^{na}\sqrt{\delta_n(1-\delta_n)}}{r} & 1 - \frac{1 + e^{na} - 2e^{na}\delta_n}{2r} \end{bmatrix},$$

$$E_2 = \frac{1}{2} \begin{bmatrix} 1 - \frac{1 + e^{na} - 2e^{na}\delta_n}{2r} & \frac{e^{na}\sqrt{\delta_n(1-\delta_n)}}{r} \\ \frac{e^{na}\sqrt{\delta_n(1-\delta_n)}}{r} & 1 + \frac{1 + e^{na} - 2e^{na}\delta_n}{2r} \end{bmatrix},$$

with

$$r \stackrel{\text{def}}{=} \frac{\sqrt{(1 + e^{na})^2 - 4e^{na}\delta_n}}{2}.$$

This leads to $\{\rho_n - e^{na}\sigma_n > 0\} = E_1$ and

$$\text{Tr}(\rho_n \{\rho_n - e^{na}\sigma_n > 0\}) = \frac{1}{2} + \frac{1 + e^{na} - 2e^{na}\delta_n}{2\sqrt{(1 + e^{na})^2 - 4e^{na}\delta_n}}.$$

Thus we have for every $a > 0$

$$\begin{aligned} \lim_{n \rightarrow \infty} \text{Tr}(\rho_n \{\rho_n - e^{na}\sigma_n > 0\}) &= 1 &\iff \lim_{n \rightarrow \infty} \delta_n &= 0, \\ \lim_{n \rightarrow \infty} \text{Tr}(\rho_n \{\rho_n - e^{na}\sigma_n > 0\}) &= 0 &\iff \lim_{n \rightarrow \infty} \delta_n &= 1, \end{aligned}$$

which yields

$$\underline{D}(\vec{\rho} \parallel \vec{\sigma}) = \begin{cases} \infty & \text{if } \lim_{n \rightarrow \infty} \delta_n = 0, \\ 0 & \text{otherwise,} \end{cases}$$

$$\overline{D}(\vec{\rho} \parallel \vec{\sigma}) = \begin{cases} 0 & \text{if } \lim_{n \rightarrow \infty} \delta_n = 1, \\ \infty & \text{otherwise.} \end{cases}$$

Furthermore, it is not difficult to see that for every $a > 0$

$$\underline{\eta}(a) = -\limsup_{n \rightarrow \infty} \frac{1}{n} \log \delta_n,$$

and letting c denote this constant $\underline{\eta}(a)$, we have

$$B_c(r \mid \vec{\rho} \parallel \vec{\sigma}) = \begin{cases} \infty & \text{if } r \leq c, \\ c & \text{if } r > c. \end{cases} \quad (49)$$

Actually, the test $T_n^{(1)} \stackrel{\text{def}}{=} \rho_n = |\psi_n\rangle\langle\psi_n|$ satisfies $\alpha_n[T_n^{(1)}] = 0$ and $\beta_n[T_n^{(1)}] = \delta_n$ for all n and hence $\underline{\eta}[\vec{T}^{(1)}] = \infty$ and $\underline{\zeta}[\vec{T}^{(1)}] = c$, while the test $T_n^{(2)} \stackrel{\text{def}}{=} \sigma_n = |\varphi_n\rangle\langle\varphi_n|$ satisfies $\eta[\vec{T}^{(2)}] = c$ and $\underline{\zeta}[\vec{T}^{(2)}] = \infty$. Equation (49) means that it suffices to consider only these extreme tests when our concern is limited to the exponents of the error probabilities. In the i.i.d. case where $\mathcal{H}_n = \mathcal{H}^{\otimes n}$, $\psi_n = \psi^{\otimes n}$ and $\varphi_n = \varphi^{\otimes n}$ for distinct unit vectors ψ and φ , we have $\underline{D}(\vec{\rho} \parallel \vec{\sigma}) = \overline{D}(\vec{\rho} \parallel \vec{\sigma}) = \infty$, which is also seen from $D(\rho \parallel \sigma) = \infty$ together with the argument of section V, and $c = -\log |\langle\psi|\varphi\rangle|^2$.

Remark 8: In the classical i.i.d. case, it follows from Sanov's theorem and Cramér's theorem in large deviation theory that, for $-D(\sigma \parallel \rho) \leq \forall a \leq D(\rho \parallel \sigma)$,

$$\underline{\eta}(a) = \overline{\eta}(a) = \min_{\tau : \tau[\log \rho - \log \sigma] \leq a} D(\tau \parallel \rho) = \max_{\theta \in \mathbb{R}} (\theta a - \psi(\theta)),$$

where $\psi(\theta) \stackrel{\text{def}}{=} \log \sum_x \rho(x)^{1+\theta} \sigma(x)^{-\theta}$, and

$$\underline{\zeta}(a) = \overline{\zeta}(a) = \min_{\tau : \tau[\log \rho - \log \sigma] \geq a} D(\tau \parallel \sigma) = a + \underline{\eta}(a).$$

Applying these relations to Theorem 3 with some additional calculations, we can derive several single-letterized expressions⁵ for $B_e(r \mid \vec{\rho} \parallel \vec{\sigma})$ (see [3], [4], [11], [13]), among which are

$$\begin{aligned} B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) &= \min_{\tau: D(\tau \parallel \rho) \leq r} D(\tau \parallel \sigma) \\ &= \max_{-1 \leq \theta < 0} \frac{(1 + \theta)r + \psi(\theta)}{\theta}. \end{aligned} \quad (50)$$

In the quantum i.i.d. case, on the other hand, we have no explicit formulas for $\underline{\eta}(a)$, $\underline{\zeta}(a)$ and $B_e(r \mid \vec{\rho} \parallel \vec{\sigma})$ at present; see [18] and [20] (section 3.4) for some partial results⁶. The mathematical difficulty arising in the study of $B_e(r \mid \vec{\rho} \parallel \vec{\sigma})$ is closely related to the absence of a “quantum large deviation theorem” applicable to $\underline{\eta}(a)$ and $\underline{\zeta}(a)$ (cf. Remark 4).

VII. EXPONENTS OF PROBABILITY OF CORRECT TESTING

Suppose that $\{\sigma_n\}$ are states (i.e., $\sigma_n[I] = 1$) and let r be a real number greater than $B^\dagger(\vec{\sigma} \parallel \vec{\rho}) = \overline{D}(\vec{\sigma} \parallel \vec{\rho})$. When the first kind error probability $\alpha_n[T_n]$ of a sequence of tests $\vec{T}$ tends to 0 with a speed not slower than e^{-nr} , the second kind error probability $\beta_n[T_n]$ inevitably tends to 1. In this case, the speed at which the probability of correct testing $1 - \beta_n[T_n]$ tends to 0 can be regarded as a measure to evaluate “badness” of $\{\sigma_n\}$. Hence, it is meaningful to investigate the slowest convergence rate of $1 - \beta_n[T_n]$ when $\alpha_n[T_n]$ is required to tend to 0 with $\alpha_n[T_n] \approx e^{-nr}$ or faster. We are thus led to introduce the following quantity:

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \stackrel{\text{def}}{=} \inf_{\vec{T}} \{ \bar{\zeta}^c[\vec{T}] \mid \underline{\eta}[\vec{T}] \geq r \}, \quad (51)$$

where

$$\bar{\zeta}^c[\vec{T}] = \limsup_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log(1 - \beta_n[T_n]) \right\}.$$

Note that $B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma})$ is defined for every r but is meaningless when $r < B^\dagger(\vec{\sigma} \parallel \vec{\rho}) = \overline{D}(\vec{\sigma} \parallel \vec{\rho})$ since it vanishes for such an r . Han [3], [4] introduced $B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma})$ for the classical hypothesis testing problem and characterized it as

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) = \inf_a \left\{ a + \eta(a) + [r - \eta(a)]_+ \right\}, \quad (52)$$

where $[t]_+ \stackrel{\text{def}}{=} \max\{t, 0\}$, assuming the two conditions that the limit $\eta(a) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \eta_n(a)$ exists for all a and that for any M there exists a K such that

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sigma_n \left[\left\{ \frac{1}{n} \log \frac{\sigma_n}{\rho_n} \geq K \right\} \right] \geq M,$$

or equivalently

$$\underline{\zeta}^c(-\infty) = \infty. \quad (53)$$

In this section we provide $B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma})$ with a new characterization which needs no extra condition. Having in mind both applicability to source coding problems and consistency with the notation in [3], [4], we exchange the roles of $\vec{\rho}$ and $\vec{\sigma}$ assumed in (2) and (3), so that

$$0 < \rho_n[I] \leq \infty \quad \text{and} \quad \sigma_n[I] = 1 \quad (54)$$

are now assumed. Accordingly, the definition of $\alpha_n[T_n]$ in (1) is changed into $\alpha_n[T_n] \stackrel{\text{def}}{=} \rho_n[I - T_n]$ together with those of $\eta_n[T_n]$, $\eta_n(a)$, $\underline{\eta}[\vec{T}]$ and $\underline{\eta}(a)$. The arguments below are based on the inequalities (15) and (16), which do not suffer from this change.

Lemma 2: For any real number a and any sequence of tests $\vec{T}$ we have

$$\bar{\zeta}^c[\vec{T}] \geq \min \{ \bar{\zeta}^c(a), a + \underline{\eta}[\vec{T}] \}. \quad (55)$$

In particular, for any $a < b$

$$\bar{\zeta}^c(b) \geq a + \underline{\eta}(b) \quad \text{if} \quad \bar{\zeta}^c(a) > \bar{\zeta}^c(b). \quad (56)$$

Proof: We have

$$\begin{aligned} \bar{\zeta}^c[\vec{T}] &= \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \beta_n^c[T_n] \\ &\geq \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \left(\beta_n^c(a) + e^{-na} \alpha_n[T_n] \right) \\ &\geq \min \left\{ \bar{\zeta}^c(a), a + \underline{\eta}[\vec{T}] \right\}, \end{aligned}$$

⁵The expressions (the first one in (50) in particular) are often referred to as *Hoeffding's theorem* after [24].

⁶Note that $B(r \mid \rho \parallel \sigma)$ in [20] corresponds to our $B_e(r \mid \vec{\sigma} \parallel \vec{\rho})$.

where the first inequality follows from (16) and the second inequality follows from the general formula (cf. (43)):

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \frac{1}{n} \log(x_n + y_n) \\ & \leq \max \left\{ \liminf_{n \rightarrow \infty} \frac{1}{n} \log x_n, \limsup_{n \rightarrow \infty} \frac{1}{n} \log y_n \right\}. \end{aligned} \quad (57)$$

■

Theorem 4: For any real number r , we have

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) = B_{e,1}^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \quad (58)$$

$$= \sup_a \min \{ \bar{\zeta}^c(a), r + a \} \quad (59)$$

$$= \inf_a \max \{ \bar{\zeta}^c(a), r + a \} \quad (60)$$

$$= r + a_0^*, \quad (61)$$

where

$$B_{e,1}^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \stackrel{\text{def}}{=} \inf_{\vec{T}} \{ \bar{\zeta}^c[\vec{T}] \mid \eta_n[T_n] \geq r \ (\forall n) \}, \quad (62)$$

$$a_0^* \stackrel{\text{def}}{=} \sup \{ a \mid \bar{\zeta}^c(a) - a \geq r \} = \inf \{ a \mid \bar{\zeta}^c(a) - a \leq r \}. \quad (63)$$

Remark 9: Note that $\geq$ and $\leq$ in the definition of a_0^* can be replaced with $>$ and $<$, respectively, because $\bar{\zeta}^c(a) - a$ is a strictly decreasing function.

Proof: Suppose that a sequence of tests $\vec{T} = \{T_n\}$ satisfies $\underline{\eta}[\vec{T}] \geq r$. It then follows from (55) that

$$\begin{aligned} \bar{\zeta}^c[\vec{T}] & \geq \sup_a \min \{ \bar{\zeta}^c(a), \underline{\eta}[\vec{T}] + a \} \\ & \geq \sup_a \min \{ \bar{\zeta}^c(a), r + a \} \\ & \geq \sup \{ r + a \mid \bar{\zeta}^c(a) \geq r + a \} = r + a_0^*, \end{aligned}$$

which proves

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \geq \sup_a \min \{ \bar{\zeta}^c(a), r + a \} \geq r + a_0^*. \quad (64)$$

Next, we show that, for arbitrarily given r , a and n , there exists a test T_n satisfying

$$\eta_n[T_n] \geq r \quad \text{and} \quad \zeta_n^c[T_n] \leq \max \{ \zeta_n^c(a), r + a \}. \quad (65)$$

When $\eta_n(a) \geq r$, it is obvious that $T_n \stackrel{\text{def}}{=} S_n(a)$ satisfies this condition. When $\eta_n(a) < r$, let

$$\begin{aligned} T_n & \stackrel{\text{def}}{=} I - e^{-n(r-\eta_n(a))} S_n^c(a) \\ & = S_n(a) + \{1 - e^{-n(r-\eta_n(a))}\} S_n^c(a). \end{aligned} \quad (66)$$

In other words, T_n is a randomized test which rejects the hypothesis ρ_n with probability $e^{-n(r-\eta_n(a))}$ when, and only when, the test $S_n(a)$ rejects ρ_n . Then we have

$$\begin{aligned} \alpha_n[T_n] & = \rho_n[e^{-n(r-\eta_n(a))} S_n^c(a)] \\ & = e^{-n(r-\eta_n(a))} \alpha_n(a) = e^{-nr} \end{aligned}$$

and

$$\begin{aligned} \beta_n^c[T_n] & = \sigma_n[e^{-n(r-\eta_n(a))} S_n^c(a)] \\ & = e^{-n(r-\eta_n(a))} \beta_n^c(a) \geq e^{-n(r+a)}, \end{aligned}$$

where the last inequality follows from (15). These are rewritten as $\eta_n[T_n] = r$ and $\zeta_n^c[T_n] \leq r + a$, and imply (65).

We have thus shown that for every r and a there exists a sequence of tests $\vec{T} = \{T_n\}$ such that $\eta_n[T_n] \geq r$ for every n and

$$\begin{aligned} \bar{\zeta}^c[\vec{T}] & \leq \limsup_{n \rightarrow \infty} \max \{ \zeta_n^c(a), r + a \} \\ & = \max \{ \bar{\zeta}^c(a), r + a \}, \end{aligned}$$

which leads to

$$\begin{aligned} B_{e,1}^*(r \mid \vec{\rho} \parallel \vec{\sigma}) &\leq \inf_a \max \{ \bar{\zeta}^c(a), r + a \} \\ &\leq \inf_a \{ r + a \mid \bar{\zeta}^c(a) \leq r + a \} = r + a_0^*. \end{aligned} \quad (67)$$

Since $B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \leq B_{e,1}^*(r \mid \vec{\rho} \parallel \vec{\sigma})$ is obvious, this together with (64) completes the proof. $\blacksquare$

Remark 10: The proof of lemma 2 can be modified, using (43) instead of (57), to yield

$$\underline{\zeta}^c[\vec{T}] \geq \min \{ \underline{\zeta}^c(a), a + \underline{\eta}[\vec{T}] \}, \quad (68)$$

whereby we can similarly show that

$$\hat{B}_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) = \hat{B}_{e,1}^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \quad (69)$$

$$= \sup_a \min \{ \underline{\zeta}^c(a), r + a \} \quad (70)$$

$$= \inf_a \max \{ \underline{\zeta}^c(a), r + a \} \quad (71)$$

$$= r + \hat{a}_0^*, \quad (72)$$

where

$$\hat{B}_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \stackrel{\text{def}}{=} \inf_{\vec{T}} \{ \underline{\zeta}^c[\vec{T}] \mid \underline{\eta}[\vec{T}] \geq r \}, \quad (73)$$

$$\hat{B}_{e,1}^*(r \mid \vec{\rho} \parallel \vec{\sigma}) \stackrel{\text{def}}{=} \inf_{\vec{T}} \{ \underline{\zeta}^c[\vec{T}] \mid \eta_n[T_n] \geq r \ (\forall n) \}, \quad (74)$$

and

$$\hat{a}_0^* \stackrel{\text{def}}{=} \sup \{ a \mid \underline{\zeta}^c(a) - a \geq r \} = \inf \{ a \mid \underline{\zeta}^c(a) - a \leq r \}. \quad (75)$$

Now let us demonstrate how Han's formula (52) is derived from Theorem 4.

Lemma 3: It always holds that

$$\bar{\zeta}^c(a) \leq a + \bar{\eta}(a) \quad (76)$$

for any a , and if a is a decreasing point of $\bar{\zeta}^c$ in the sense that $\bar{\zeta}^c(a - \varepsilon) > \bar{\zeta}^c(a + \varepsilon)$ for any $\varepsilon > 0$, then we also have

$$\bar{\zeta}^c(a + 0) \geq a + \underline{\eta}(a + 0). \quad (77)$$

Proof: The inequalities are immediate from (15) and (56), respectively. $\blacksquare$

Corollary 1: It always holds that

$$\begin{aligned} B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) &\leq \inf_a \left(a + \max \{ \bar{\eta}(a), r \} \right) \\ &= \inf_a \left\{ a + \bar{\eta}(a) + [r - \bar{\eta}(a)]_+ \right\}, \end{aligned} \quad (78)$$

while we have

$$\begin{aligned} B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) &\geq \inf_a \left(a + \max \{ \underline{\eta}(a), r \} \right) \\ &= \inf_a \left\{ a + \underline{\eta}(a) + [r - \underline{\eta}(a)]_+ \right\} \end{aligned} \quad (79)$$

if

$$r \leq \bar{\zeta}^c(-\infty) - \sup \{ a \mid \bar{\zeta}^c(-\infty) = \bar{\zeta}^c(a) \}. \quad (80)$$

In particular, if the limit $\eta(a) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \eta_n(a)$ exists for all a and if

$$\bar{\zeta}^c(-\infty) = \infty \quad \text{or} \quad \{ a \mid \bar{\zeta}^c(-\infty) = \bar{\zeta}^c(a) \} = \phi, \quad (81)$$

then Han's formula (52) is valid for all r .

Proof: Since the first inequality is immediate from Theorem 4 and (76), we only prove the second one. Invoking Theorem 4 again, it suffices to show that, for any r satisfying (80),

$$\inf_a \max \{ \bar{\zeta}^c(a), r + a \} \geq \inf_a \left(a + \max \{ \underline{\eta}(a), r \} \right). \quad (82)$$

Define

$$b_0 \stackrel{\text{def}}{=} \sup \{ a \mid \bar{\zeta}^c(-\infty) = \bar{\zeta}^c(a) \},$$

including the case when $\{ a \mid \bar{\zeta}^c(-\infty) = \bar{\zeta}^c(a) \} = \phi$ and $b_0 = -\infty$, and let a be an arbitrary number satisfying $a > b_0$. Then we have $\bar{\zeta}^c(-\infty) > \bar{\zeta}^c(a)$, which means that there exists a number b such that $b < a$ and $\bar{\zeta}^c(b) > \bar{\zeta}^c(a)$. The supremum of

such numbers b , denoted by $\bar{b} = \bar{b}(a)$, satisfies $\bar{b} \leq a$ and $\bar{\zeta}^c(\bar{b} - \varepsilon) > \bar{\zeta}^c(a) \geq \bar{\zeta}^c(\bar{b} + \varepsilon)$ for every $\varepsilon > 0$, which implies that $\bar{b}$ is a decreasing point of $\underline{\zeta}^c$ and $\bar{\zeta}^c(a) \geq \bar{\zeta}^c(\bar{b} + 0)$. Hence we have

$$\begin{aligned} \max \{\bar{\zeta}^c(a), r + a\} &\geq \max \{\bar{\zeta}^c(\bar{b} + 0), r + \bar{b}\} \\ &\geq \max \{\underline{\eta}(\bar{b} + 0) + \bar{b}, r + \bar{b}\} \\ &= \lim_{\varepsilon \downarrow 0} \left(\bar{b} + \varepsilon + \max \{\underline{\eta}(\bar{b} + \varepsilon), r\} \right) \\ &\geq \inf_b \left(b + \max \{\underline{\eta}(b), r\} \right), \end{aligned}$$

where the second inequality follows from (77), and therefore

$$\inf_{a: a > b_0} \max \{\bar{\zeta}^c(a), r + a\} \geq \inf_a \left(a + \max \{\underline{\eta}(a), r\} \right). \quad (83)$$

This proves (82) when $b_0 = -\infty$. In the case when $b_0 > -\infty$, we have

$$\begin{aligned} \inf_{a: a < b_0} \max \{\bar{\zeta}^c(a), r + a\} &\geq \inf_{a: a < b_0} \bar{\zeta}^c(a) \\ &= \bar{\zeta}^c(-\infty) \\ &\geq \max \{\bar{\zeta}^c(b_0), r + b_0\}, \end{aligned} \quad (84)$$

where the last inequality follows from the nonincreasing property of $\bar{\zeta}^c$ and (80), and in addition we have

$$\begin{aligned} \max \{\bar{\zeta}^c(b_0), r + b_0\} &\geq \max \{\bar{\zeta}^c(b_0 + 0), r + b_0\} \\ &= \lim_{\varepsilon \downarrow 0} \max \{\bar{\zeta}^c(b_0 + \varepsilon), r + b_0 + \varepsilon\} \\ &\geq \inf_{a: a > b_0} \max \{\bar{\zeta}^c(a), r + a\}. \end{aligned} \quad (85)$$

Now the desired inequality (82) follows from (83), (84) and (85). $\blacksquare$

Remark 11: The first condition in (81) in terms of $\bar{\zeta}^c$ is weaker than the original condition (53) in terms of $\underline{\zeta}^c$. The second condition in (81) means that for any number a there always exists a number $b < a$ such that $\bar{\zeta}^c(b) > \bar{\zeta}^c(a)$. The second is not implied by the first, since there may be a number a such that $\bar{\zeta}^c(b) = \infty$ for all $b > a$.

Remark 12: It is easy to see that if the condition (80) is not satisfied then

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) = \underline{\zeta}^c(-\infty).$$

Example 3: Consider the following classical hypothesis testing problem: $\mathcal{X}^n = \{x_0, x_1\}$ for all n on which probability distributions ρ_n and σ_n are defined by

$$\rho_n(x_0) = e^{-nb_n}, \quad \rho_n(x_1) = 1 - e^{-nb_n}, \quad (86)$$

$$\sigma_n(x_0) = e^{-nc}, \quad \sigma_n(x_1) = 1 - e^{-nc}, \quad (87)$$

where b_n is a positive sequence obeying $\lim_{n \rightarrow \infty} b_n = \infty$ and c is a positive constant. Then the limits $\eta(a) = \lim_{n \rightarrow \infty} \eta_n(a)$ and $\zeta^c(a) = \lim_{n \rightarrow \infty} \zeta_n^c(a)$ exist for all a and satisfy

$$\eta(a) = \begin{cases} 0 & \text{if } a > 0 \\ \infty & \text{if } a \leq 0 \end{cases} \quad \text{and} \quad \zeta^c(a) = \begin{cases} 0 & \text{if } a > 0 \\ c & \text{if } a \leq 0, \end{cases}$$

where we have chosen $S_n(a) = \{\rho_n - e^{na}\sigma_n > 0\}$ in (6). Note also that $\underline{D}(\vec{\sigma} \parallel \vec{\rho}) = \overline{D}(\vec{\sigma} \parallel \vec{\rho}) = 0$. It is then immediate from Theorem 4 that

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) = \begin{cases} c & \text{if } r \geq c \\ r & \text{if } 0 \leq r \leq c \\ 0 & \text{if } r \leq 0, \end{cases}$$

while we have

$$\inf_a \left\{ a + \eta(a) + [r - \eta(a)]_+ \right\} = \begin{cases} r & \text{if } r \geq 0 \\ 0 & \text{if } r \leq 0. \end{cases}$$

Since $\zeta^c(-\infty) = c$ and $\sup \{a \mid \zeta^c(-\infty) = \zeta^c(a)\} = 0$, the condition (80) for validity of Han's formula turns out to be $r \leq c$, which just explains the above situation.

Remark 13: In the classical i.i.d. case, Han and Kobayashi [25] (see also [26]) obtained a compact expression for $B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma})$ in the form

$$B_e^*(r \mid \vec{\rho} \parallel \vec{\sigma}) = \min_{\tau: D(\tau \parallel \rho) \leq r} \{D(\tau \parallel \sigma) + r - D(\tau \parallel \rho)\} \quad (88)$$

with noting that the RHS can be represented as $\min_{\tau: D(\tau \| \rho) \geq r} D(\tau \| \sigma)$ when r is sufficiently near $D(\sigma \| \rho)$. We also have an expression in the form⁷

$$B_e^*(r | \vec{\rho} \| \vec{\sigma}) = \max_{\theta \leq -1} \frac{(1 + \theta)r + \psi(\theta)}{\theta}, \quad (89)$$

where $\psi(\theta) = \log \sum_x \rho(x)^{1+\theta} \sigma(x)^{-\theta}$ is the same function as defined in Remark 8. These expressions can be derived by applying large deviation theorems to Theorem 4 (or to (52) as in [3], [4]) (cf. Remark 8). For the quantum i.i.d. case, it was shown in [21] that inequality (33), with ρ and σ exchanged, yields a lower bound on $B_e^*(r | \vec{\rho} \| \vec{\sigma})$ in the same form as the RHS of (89) except that the range of $\max$ is restricted to $-2 \leq \theta \leq -1$ (; see [22] for a simple derivation). This restriction has been relaxed to $\theta \leq -1$ just as (89) by [27] and [20] (section 3.4)⁸. Some further results on $B_e^*(r | \vec{\rho} \| \vec{\sigma})$ are also found in [20] including a quantum extension of (89) (not a bound but an identity) in terms of a variant of $\psi(\theta)$ which is defined in a limiting form⁹.

Before concluding this section, we introduce the dual of $B_e^*(r | \vec{\rho} \| \vec{\sigma})$ by

$$\begin{aligned} B_e^{**}(r | \vec{\rho} \| \vec{\sigma}) &\stackrel{\text{def}}{=} \sup_{\vec{T}} \{ \eta[\vec{T}] | \bar{\zeta}^c[\vec{T}] \leq r \} \\ &= \sup \{ r' | B_e^*(r' | \vec{\rho} \| \vec{\sigma}) \leq r \} \end{aligned} \quad (90)$$

and provide this with a general characterization, which will be applied to the source coding problem in the next section.

Theorem 5: We have

$$B_e^{**}(r | \vec{\rho} \| \vec{\sigma}) = r - a_0^{**}, \quad (91)$$

where

$$a_0^{**} \stackrel{\text{def}}{=} \inf \{ a | \bar{\zeta}^c(a) \leq r \} = \sup \{ a | \bar{\zeta}^c(a) > r \}. \quad (92)$$

Proof: Obvious from the following equivalence:

$$\begin{aligned} B_e^*(r' | \vec{\rho} \| \vec{\sigma}) \leq r &\Leftrightarrow \sup_a \min \{ \bar{\zeta}^c(a), r' + a \} \leq r \\ &\Leftrightarrow \forall a \left(a > r - r' \Rightarrow \bar{\zeta}^c(a) \leq r \right) \\ &\Leftrightarrow r' \leq r - a_0^{**}. \end{aligned}$$

■

VIII. APPLICATION TO CLASSICAL FIXED-LENGTH SOURCE CODING

Let $\vec{\rho} = \{\rho_n\}_{n=1}^\infty$ be a classical general source; i.e., let ρ_n be a probability distribution P_{X^n} on a finite or countably infinite set $\mathcal{X}^n$ for each n . A (possibly stochastic) fixed-length coding system for this source is generally represented by a sequence $\vec{\Phi} = \{\Phi_n\}_{n=1}^\infty$ of $\Phi_n = (\mathcal{Y}_n, F_n, G_n)$, where $\mathcal{Y}_n$ is a finite set, $F_n = \{F_n(y | x)\}$ is a channel from $\mathcal{X}^n$ to $\mathcal{Y}_n$ representing an encoder, and $G_n = \{G_n(x | y)\}$ is a channel from $\mathcal{Y}_n$ to $\mathcal{X}^n$ representing a decoder. The size and the error probability of Φ_n are respectively defined by $|\Phi_n| \stackrel{\text{def}}{=} |\mathcal{Y}_n|$ and

$$\gamma_n[\Phi_n] \stackrel{\text{def}}{=} 1 - \sum_{x \in \mathcal{X}^n} \sum_{y \in \mathcal{Y}_n} \rho_n(x) F_n(y | x) G_n(x | y).$$

Now, let σ_n be the counting measure on $\mathcal{X}^n$; i.e., $\sigma_n[T_n] = \sum_{x \in \mathcal{X}^n} T_n(x)$. Then the source coding problem for ρ_n can be reduced to the generalized hypothesis testing problem for $\{\rho_n, \sigma_n\}$ as follows. For an arbitrary coding system Φ_n , a test T_n is defined by

$$T_n(x) = \sum_{y \in \mathcal{Y}_n} F_n(y | x) G_n(x | y), \quad (93)$$

which satisfies

$$\gamma_n[\Phi_n] = 1 - \rho_n[T_n] = \alpha_n[T_n] \quad (94)$$

and

$$\begin{aligned} |\Phi_n| = |\mathcal{Y}_n| &= \sum_{y \in \mathcal{Y}_n} 1 \geq \sum_{y \in \mathcal{Y}_n} \sum_{x \in \mathcal{X}^n} F_n(y | x) G_n(x | y) \\ &= \sigma_n[T_n] = \beta_n[T_n]. \end{aligned} \quad (95)$$

⁷To the authors' knowledge, this type of expression for B_e^* first appeared in [21] even for the classical case.

⁸Note that $B^*(r | \rho \| \sigma)$ in [20] corresponds to our $B_e^*(r | \vec{\sigma} \| \vec{\rho})$.

⁹This result has not appeared in the original Japanese edition of [20].

Conversely, for an arbitrary deterministic (i.e., $\{0, 1\}$ -valued) test T_n we can construct a coding system $\Phi_n = (\mathcal{Y}_n, F_n, G_n)$ satisfying $\gamma_n[\Phi_n] = \alpha_n[T_n]$ and $|\Phi_n| = \beta_n[T_n]$ by setting $\mathcal{Y}_n = \{x \in \mathcal{X}^n \mid T_n(x) = 1\}$ and $F_n(y \mid x) = G_n(x \mid y) = 1$ if $y = x \in \mathcal{Y}_n$. Noting that the direct (achievability) parts of Theorems 1 and 3 have been shown by using only deterministic tests (by setting $S_n(a)$ to be $\{\rho_n - e^{na}\sigma_n > 0\}$ or $\{\rho_n - e^{na}\sigma_n \geq 0\}$) and that $\limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| = -\underline{\zeta}[\vec{T}]$ when $|\Phi_n| = \beta_n[T_n]$, we immediately obtain the following identities:

Theorem 6: We have

$$\begin{aligned} R(\varepsilon \mid \vec{\rho}) &\stackrel{\text{def}}{=} \inf_{\vec{\Phi}} \{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| \mid \limsup_{n \rightarrow \infty} \gamma_n[\Phi_n] \leq \varepsilon \} \\ &= -B(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) = -\underline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) \end{aligned} \quad (96)$$

$$= \inf \{ a \mid \limsup_{n \rightarrow \infty} P_{X^n} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \geq a \right\} \leq \varepsilon \}, \quad (97)$$

$$\begin{aligned} R^\dagger(\varepsilon \mid \vec{\rho}) &\stackrel{\text{def}}{=} \inf_{\vec{\Phi}} \{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| \mid \liminf_{n \rightarrow \infty} \gamma_n[\Phi_n] < \varepsilon \} \\ &= -B^\dagger(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) = -\overline{D}(\varepsilon \mid \vec{\rho} \parallel \vec{\sigma}) \end{aligned} \quad (98)$$

$$= \inf \{ a \mid \liminf_{n \rightarrow \infty} P_{X^n} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \geq a \right\} < \varepsilon \}, \quad (99)$$

$$\begin{aligned} R(\vec{\rho}) &\stackrel{\text{def}}{=} R(0 \mid \vec{\rho}) \\ &= \inf \{ R \mid \exists \vec{\Phi}, \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| \leq R \text{ and } \lim_{n \rightarrow \infty} \gamma_n[\Phi_n] = 0 \} \\ &= -B(\vec{\rho} \parallel \vec{\sigma}) = -\underline{D}(\vec{\rho} \parallel \vec{\sigma}) \end{aligned} \quad (100)$$

$$= \overline{H}(\vec{\rho}) \stackrel{\text{def}}{=} \text{p-lim sup}_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \right\}, \quad (101)$$

$$\begin{aligned} R^\dagger(\vec{\rho}) &\stackrel{\text{def}}{=} R^\dagger(1 \mid \vec{\rho}) \\ &= \sup \{ R \mid \forall \vec{\Phi}, \text{ if } \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| \leq R \text{ then } \lim_{n \rightarrow \infty} \gamma_n[\Phi_n] = 1 \} \\ &= -B^\dagger(\vec{\rho} \parallel \vec{\sigma}) = -\overline{D}(\vec{\rho} \parallel \vec{\sigma}) \end{aligned} \quad (102)$$

$$= \underline{H}(\vec{\rho}) \stackrel{\text{def}}{=} \text{p-lim inf}_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \right\}, \quad (103)$$

and

$$\begin{aligned} R_e(r \mid \vec{\rho}) &\stackrel{\text{def}}{=} \inf_{\vec{\Phi}} \{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| \mid \liminf_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log \gamma_n[\Phi_n] \right\} \geq r \} \\ &= -B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) \end{aligned} \quad (104)$$

$$= \sup_a \{ a - \underline{\sigma}(a) \mid \underline{\sigma}(a) < r \}, \quad (105)$$

where

$$\begin{aligned} \underline{\sigma}(a) &\stackrel{\text{def}}{=} \underline{\eta}(-a) \\ &= \liminf_{n \rightarrow \infty} -\frac{1}{n} \log P_{X^n} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \geq a \right\}. \end{aligned} \quad (106)$$

Remark 14: $R(\vec{\rho})$ is the optimal compression rate with asymptotically vanishing error probability and Equation (101), which was originally shown in [1], means that it always equals the spectral sup-entropy rate $\overline{H}(\vec{\rho})$. The source $\vec{\rho}$ is said to have the strong converse property when $\limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| < R(\vec{\rho})$ implies $\lim_{n \rightarrow \infty} \gamma_n[\Phi_n] = 1$, or equivalently when $R(\vec{\rho}) = R^\dagger(\vec{\rho})$. As was pointed out in [1], this property is equivalent to $\overline{H}(\vec{\rho}) = \underline{H}(\vec{\rho})$, which is now obvious from (101) and (103). Equation (97) is found in [28], and (105) in [5]. Although the use of the symbol σ for different notions, for the counting measures and for the function (106), may be a little confusing, it will be helpful for comparing our results with those of [5].

Next, let us turn to the following quantity:

$$\begin{aligned} R_e^*(r \mid \vec{\rho}) &\stackrel{\text{def}}{=} \inf_{\vec{\Phi}} \{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| \mid \limsup_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log(1 - \gamma_n[\Phi_n]) \right\} \leq r \}. \end{aligned} \quad (107)$$

Han [3], [5] proved that for any $r \geq 0$

$$R_e^*(r | \vec{\rho}) = \inf \{h \geq 0 \mid \inf_a \{\sigma^*(a) + [a - \sigma^*(a) - h]^+\} \leq r\} \quad (108)$$

under the assumption that the following limit exists for all a :

$$\sigma^*(a) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} -\frac{1}{n} \log P_{X^n} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \leq a \right\}.$$

A general formula for $R_e^*(r | \vec{\rho})$ which needs no additional assumption is given below.

Theorem 7: For any $r \geq 0$ we have

$$R_e^*(r | \vec{\rho}) = \max \{b_0 - r, 0\}, \quad (109)$$

where

$$b_0 \stackrel{\text{def}}{=} \sup \{a \mid \bar{\sigma}^*(a) > r\} = \inf \{a \mid \bar{\sigma}^*(a) \leq r\}$$

and

$$\bar{\sigma}^*(a) \stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} -\frac{1}{n} \log P_{X^n} \left\{ -\frac{1}{n} \log P_{X^n}(X^n) \leq a \right\}.$$

Proof: Since

$$\bar{\sigma}^*(a) = \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \rho_n [\{\rho_n - e^{-na} \sigma_n \geq 0\}] = \bar{\eta}^c(-a),$$

it follows from Theorem 5, with $\vec{\rho}$ and $\vec{\sigma}$ exchanged, that

$$B_e^{**}(r | \vec{\sigma} \| \vec{\rho}) = r - b_0.$$

Hence (109) is equivalent to

$$R_e^*(r | \vec{\rho}) = \max \{-B_e^{**}(r | \vec{\sigma} \| \vec{\rho}), 0\}. \quad (110)$$

Here it is easy to see LHS $\geq$ RHS from (94), (95) and

$$\begin{aligned} -B_e^{**}(r | \vec{\sigma} \| \vec{\rho}) &= -\sup_{\vec{T}} \{ \zeta[\vec{T}] \mid \bar{\eta}^c[\vec{T}] \leq r \} \\ &= \inf_{\vec{T}} \{ \limsup_{n \rightarrow \infty} \frac{1}{n} \log \beta_n[T_n] \mid \limsup_{n \rightarrow \infty} -\frac{1}{n} \log(1 - \alpha_n[T_n]) \leq r \}. \end{aligned}$$

Let us show the converse inequality LHS $\leq$ RHS. Since

$$-B_e^{**}(r | \vec{\sigma} \| \vec{\rho}) = \inf \{r' \mid B_e^*(-r' | \vec{\sigma} \| \vec{\rho}) \leq r\},$$

it is sufficient to show that

$$R_e^*(r | \vec{\rho}) \leq r' \quad \text{if} \quad B_e^*(-r' | \vec{\sigma} \| \vec{\rho}) \leq r \quad \text{and} \quad r' > 0.$$

This is equivalent to the proposition that for any $r' > 0$ there exists a sequence of codes $\vec{\Phi}$ such that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \log |\Phi_n| &\leq r', \quad \text{and} \\ \limsup_{n \rightarrow \infty} -\frac{1}{n} \log(1 - \gamma_n[T_n]) &\leq B_e^*(-r' | \vec{\sigma} \| \vec{\rho}). \end{aligned}$$

This proposition follows if the infimum of

$$B_e^*(-r' | \vec{\sigma} \| \vec{\rho}) = \inf_{\vec{T}} \{ \bar{\eta}^c[\vec{T}] \mid \zeta[\vec{T}] \geq -r' \}$$

can be attained by a sequence of deterministic tests $\vec{T}$ when $r' > 0$. Recalling the proof of Theorem 4 and applying it to the present situation, it suffices to show that for any $a \in \mathbb{R}$, $r' > 0$, $\delta > 0$ and for any sufficiently large n there exists a deterministic test satisfying

$$\zeta_n[T_n] \geq -r' - \delta \quad \text{and} \quad \eta_n^c[T_n] \leq \max \{\eta_n^c(a), -r' - a\}, \quad (111)$$

which corresponds to (65) with a slight modification. Let $S_n(a)$ be chosen to be deterministic in (6) and identify it with its acceptance region (e.g., $S_n(a) = \{x \in \mathcal{X}^n \mid \rho_n(x) - e^{na} \sigma_n(x) > 0\}$). It is then obvious that $T_n \stackrel{\text{def}}{=} S_n(a)$ satisfies (111) if $\zeta_n(a) \geq -r'$. Suppose $\zeta_n(a) < -r'$, which means $\sigma_n[S_n(a)] = |S_n(a)| > e^{nr'}$. Then there exists a subset $T_n \subset S_n(a)$, which is regarded as a deterministic test, satisfying

$$\sigma[T_n] = |T_n| = \lceil e^{nr'} \rceil \quad \text{and} \quad \rho_n[T_n] \geq \frac{\lceil e^{nr'} \rceil}{|S_n(a)|} \rho_n[S_n(a)].$$

Using (9) we have $\rho_n[T_n] \geq e^{n(a+r')}$, and T_n satisfies (111). ■

Now we can see that Han's formula (108) is very near to the true general formula. Actually, if $\sigma^*(a)$ is simply replaced with $\bar{\sigma}^*(a)$, it becomes equivalent to (109) as follows. Noting that $\bar{\sigma}^*(a)$ is monotonically nonincreasing, we have

$$\begin{aligned}
& \inf_a \{ \bar{\sigma}^*(a) + [a - \bar{\sigma}^*(a) - h]^+ \} \leq r \\
\Leftrightarrow & \inf_a \max \{ \bar{\sigma}^*(a), a - h \} \leq r \\
\Leftrightarrow & \forall \delta > 0, \exists a, \bar{\sigma}^*(a) \leq r + \delta \text{ and } a - h \leq r + \delta \\
\Leftrightarrow & h \geq \sup_{\delta > 0} \inf_a \{ a - (r + \delta) \mid \bar{\sigma}^*(a) \leq r + \delta \} \\
\Leftrightarrow & h \geq \sup_{\delta > 0} \sup_a \{ a - (r + \delta) \mid \bar{\sigma}^*(a) > r + \delta \} \\
\Leftrightarrow & h \geq \sup_a \{ a - r \mid \exists \delta > 0, \bar{\sigma}^*(a) > r + \delta \} \\
\Leftrightarrow & h \geq \sup_a \{ a - r \mid \bar{\sigma}^*(a) > r \} = b_0 - r,
\end{aligned}$$

and therefore

$$\begin{aligned}
& \inf \{ h \geq 0 \mid \inf_a \{ \sigma^*(a) + [a - \bar{\sigma}^*(a) - h]^+ \} \leq r \} \\
& = \max \{ b_0 - r, 0 \}.
\end{aligned}$$

Remark 15: Iriyama (and Ihara) [29], [30] obtained other forms of general formulas for $R_e(r \mid \vec{\rho})$ and $R_e^*(r \mid \vec{\rho})$ from a different point of view.

Remark 16: Although we have treated only classical source coding here, extension to some quantum settings is actually possible. Of the two major coding schemes proposed for the quantum pure state source coding, namely visible coding and blind coding, the former is less restrictive and hence needs in general a more careful or stronger argument than the latter when showing the converse part of a theorem concerning a limit on all possible codes. The situation is reversed when showing the direct (achievability) part. It is easy to see that the direct parts of Theorems 6 and 7 are straightforwardly extended to visible coding, and that the direct part only of the arguments concerning $R(\vec{\rho})$ and $R_e(r \mid \vec{\rho})$ in Theorems 6 is applicable to blind coding, while it is not clear whether other bounds in Theorems 6 and 7 are achievable for blind coding. On the other hand, it has been shown in [31] that the inequality

$$\gamma_n[\Phi_n] + e^{na}|\Phi_n| \geq \rho_n[\{\rho_n - e^{na}\sigma_n \leq 0\}],$$

which follows from (14), (94) and (95), can be extended to visible coding just in the same form. Since the converse parts of our theorems are direct consequences of this inequality, they are extended to visible coding, and hence to blind coding as well. We thus have the same formula as Theorem 6 for both visible and blind coding, and Theorem 7 for visible coding. See [31] for details. Hayashi [32] showed that these values $R^\dagger(\epsilon \mid \vec{\rho})$, $R^\dagger(\vec{\rho})$, $R_e(r \mid \vec{\rho})$ have other operational meaning. He also treated these values when the quantum information source is given by the thermal state of Hamiltonian with interaction. That is, using this discussion, we can treat the bounds $R^\dagger(\epsilon \mid \vec{\rho})$, $R^\dagger(\vec{\rho})$, $R_e(r \mid \vec{\rho})$ in this case.

Remark 17: Recently, Hayashi[33] clarified the relation between $R(\epsilon \mid \vec{\rho})$ and $R^\dagger(\epsilon \mid \vec{\rho})$ from a wider view point.

IX. CONCLUDING REMARKS

We have demonstrated that the information-spectrum analysis made by Han for the classical hypothesis testing for simple hypotheses, together with the fixed-length source coding, can be naturally extended to a unifying framework including both the classical and quantum generalized hypothesis testing. The generality of theorems and the simplicity of proofs have been thoroughly pursued and have yielded some improvements of the original classical results.

The significance of our results for quantum information theory is not so clear at present, since our knowledge of the asymptotic behavior of the quantum information spectrum $\rho_n[\{\rho_n - e^{na}\sigma_n > 0\}] = \text{Tr}(\rho_n\{\rho_n - e^{na}\sigma_n > 0\})$ is insufficient even for the i.i.d. case $\rho_n = \rho^{\otimes n}$, $\sigma_n = \sigma^{\otimes n}$. Therefore we cannot obtain compact and computable representations of information-spectrum quantities. Nevertheless, the fact that the asymptotic characteristics of quantum hypothesis testing are represented in terms of the information spectrum seems to suggest the importance of studying quantum information theory from the information-spectrum viewpoint. An attempt in this direction is found in [19], where a similar approach to [2] is made for the general (classical-)quantum channels. As an application the capacity formula for quantum stationary memoryless channels [34], [35] is provided with a new simple proof by linking it to the quantum Stein's lemma via our Theorem 2.

Finally, we mention some remarkable progresses in related subjects reported after submitting the accepted version of the present paper. The quantum Chernoff bound for symmetric Bayesian discrimination of two i.i.d. states has been established by [36] and [37]. Based on an inequality shown in [37], it has been proved by [38] that $B_e(r \mid \vec{\rho} \parallel \vec{\sigma})$ for the quantum i.i.d. case satisfies (cf. equation (50))

$$B_e(r \mid \vec{\rho} \parallel \vec{\sigma}) \geq \max_{-1 \leq \theta < 0} \frac{(1 + \theta)r + \psi(\theta)}{\theta},$$

where $\psi(\theta) \stackrel{\text{def}}{=} \log \text{Tr} \rho^{1+\theta} \sigma^{-\theta}$. This is the tightest lower bound on $B_e(r \|\vec{\rho} \|\vec{\sigma})$ of those obtained so far. Moreover it seems natural to conjecture that the bound achieves the equality in general.

REFERENCES

- [1] T.S. Han and S. Verdú, "Approximation theory of output statistics," *IEEE Trans. Inform. Theory*, vol.39, no.3, 752–772, 1993.
- [2] S. Verdú and T.S. Han, "A general formula for channel capacity," *IEEE Trans. Inform. Theory*, vol.40, no.4, 1147–1157, 1994.
- [3] T.S. Han, *Information-Spectrum Methods in Information Theory*, Springer, 2003; the original Japanese edition was published from Baifukan-Press, Tokyo, in 1998.
- [4] T.S. Han, "Hypothesis testing with the general source," *IEEE Trans. Inform. Theory*, vol.46, no.7, 2415–2427, 2000.
- [5] T.S. Han, "The reliability functions of the general source with fixed-length coding," *IEEE Trans. Inform. Theory*, vol.46, no.6, 2117–2132, 2000.
- [6] T.S. Han, "An information-spectrum approach to source coding theorems with a fidelity criterion," *IEEE Trans. Inform. Theory*, vol.43, no.4, 1145–1164, 1997.
- [7] A. S. Holevo, "An analog of the theory of statistical decisions in noncommutative theory of probability," *Trudy Moskov. Mat. Obšč.*, vol. 26, 133–149 1972. (English translation is *Trans. Moscow Math. Soc.*, vol. 26, 133–149 1972.)
- [8] C.W. Helstrom, *Quantum Detection and Estimation Theory*, Academic Press, New York, 1976.
- [9] P.N. Chen, "General formulas for the Neyman-Pearson type-II error exponent subject to fixed and exponential type-I error bounds," *IEEE Trans. Inform. Theory*, vol.42, no.1, 316–323, 1996.
- [10] S. Verdú: private communication to T.S. Han, 1994.
- [11] R.E. Blahut, *Principles and Practice of Information Theory*, Addison-Wesley, 1987.
- [12] T.M. Cover and J.A. Thomas, *Elements of Information Theory*, Wiley, 1991; 2nd edition, 2006.
- [13] A. Dembo and O. Zeitouni, *Large Deviation Techniques and Applications*, Jones and Bartlett, 1993; 2nd ed., Springer, 1998.
- [14] M. Ohya and D. Petz, *Quantum Entropy and Its Use*, Springer-Verlag, 1993; 2nd ed., Springer, 2004.
- [15] F. Hiai and D. Petz, "The proper formula for relative entropy and its asymptotics in quantum probability," *Commun. Math. Phys.*, vol.143, 99–114, 1991.
- [16] M. Hayashi, "Asymptotics of quantum relative entropy from a representation theoretical viewpoint," *J. Phys. A: Math. Gen.*, vol.34, 3413–3419, 2001.
- [17] M. Hayashi, "Optimal sequences of POVMs in the sense of Stein's lemma in quantum hypothesis testing," *J. Phys. A: Math. and Gen.*, vol.35, 10759–10773, 2002.
- [18] T. Ogawa and M. Hayashi, "On error exponents in quantum hypothesis testing," *IEEE Trans. Inform. Theory*, vol.50, no.6, 1368–1372, 2004.
- [19] M. Hayashi and H. Nagaoka, "General formulas for capacity of classical-quantum channels," *IEEE Trans. Inform. Theory*, vol.49, 1753–1768, 2003.
- [20] M. Hayashi, *Quantum Information: An Introduction*, Springer, 2006 (; the original Japanese edition was published from Saiensu-sha, Tokyo, 2004).
- [21] T. Ogawa and H. Nagaoka, "Strong converse and Stein's lemma in quantum hypothesis testing," *IEEE Trans. Inform. Theory*, vol.46, no.7, 2428–2433, 2000.
- [22] H. Nagaoka, "Strong converse theorems in quantum information theory," Proc. ERATO Workshop on Quantum Information Science 2001, 33, 2001; appears also in M. Hayashi (ed.), *Asymptotic Theory of Quantum Statistical Inference*, World Scientific, 2005.
- [23] I. Bjelaković et al., "A quantum version of Sanov's theorem", *Commun. Math. Phys.*, vol.260, 659–671, 2005.
- [24] W. Hoeffding, "Asymptotically optimal test for multinomial distributions", *Ann. Math. Statist.*, vol.36, 369–400, 1965.
- [25] T.S. Han and K. Kobayashi, "The strong converse theorem for hypothesis testing," *IEEE Trans. Inform. Theory*, vol.35, 178–180, 1989.
- [26] K. Nakagawa and F. Kanaya, "On the converse theorem in statistical hypothesis testing," *IEEE Trans. Inform. Theory*, vol.39, 623–628, 1993.
- [27] M. Hayashi, "Quantum hypothesis testing for the quantum general hypotheses," Proc. Symp. Inform. Theory and Its Appl., 591–594, 2001 (in Japanese).
- [28] Y. Steinberg and S. Verdú, "Simulation of random processes and rate-distortion theory," *IEEE Trans. Inform. Theory*, vol.40, no.3, 634–646, 1994.
- [29] K. Iriyama, "Probability of error for the fixed-length source coding of general sources," *IEEE Trans. Inform. Theory*, vol.47, no.4, 1537–1543, 2001.
- [30] K. Iriyama and S. Ihara, "The error exponent and minimum achievable rates for the fixed-length coding of general sources," *IEICE Trans. Fundamentals*, vol.E84-A, no.10, 2466–2473, 2001.
- [31] M. Hayashi, "Exponents of quantum fixed-length pure state source coding," *Phys. Rev. A*, vol.66, 032321, 2002.
- [32] M. Hayashi, "General formulas for fixed-length quantum entanglement concentration," *IEEE Trans. Inform. Theory*, vol.52, no.5, 1904–1921, 2006.
- [33] M. Hayashi, "Second order asymptotics in fixed-length source coding and intrinsic randomness," LANL E-print cs.IT/0503089.
- [34] A.S. Holevo, "The capacity of the quantum channel with general signal states," *IEEE Trans. Inform. Theory*, vol.44, 269–273, 1998.
- [35] B. Schumacher and M.D. Westmoreland, "Sending classical information via noisy quantum channels," *Phys. Rev. A*, vol.56, 131–138, 1997.
- [36] M. Nussbaum and A. Szkola, "A lower bound of Chernoff type for symmetric quantum hypothesis testing," LANL E-print quant-ph/0607216.
- [37] K.M.R. Audenaert et al., "The Quantum Chernoff Bound," LANL E-print quant-ph/0610027.
- [38] M. Hayashi, "Error exponent in asymmetric quantum hypothesis testing and its application to classical-quantum channel coding," LANL E-print quant-ph/0611013.

PLACE
PHOTO
HERE

Hiroshi Nagaoka (M'88) was born in Tokyo in 1955. He received the B. Eng. and M. Eng. degrees in 1980 and 1982, respectively, from the University of Tokyo and the Dr. Eng. degree from Osaka University in 1987.

He worked at the Tokyo Engineering University from 1986 to 1989, at Hokkaido University from 1989 to 1993, and is currently an associate professor at the University of Electro-Communications. His research interests include information geometry, quantum information theory and quantum statistical inference.

PLACE
PHOTO
HERE

Masahito Hayashi was born in Japan in 1971. He received the B. S. degree from Faculty of Sciences in Kyoto University, Japan, in 1994 and the M. S. and Ph. D. degrees in Mathematics from Kyoto University, Japan, in 1996 and 1999, respectively.

He worked in Kyoto University as a Research Fellow of the Japan Society of the Promotion of Science (JSPS) from 1998 to 2000, and worked in the Laboratory for Mathematical Neuroscience, Brain Science Institute, RIKEN from 2000 to 2003. In 2003, he joined Quantum Computation and Information Project, Japan Science and Technology Agency (JST) as the Research Head. He also works in Superrobust Computation Project Information Science and Technology Strategic Core (21st Century COE by MEXT) Graduate School of Information Science and Technology The University of Tokyo as Adjunct Associate Professor from 2004. He is an Editorial Board of International Journal of Quantum Information. His research interests include quantum information theory and quantum statistical inference.